JOURNAL OF FORMALIZED MATHEMATICS
Volume 9, Released 1997, Published 2003
Inst. of Computer Science, Univ. of Białystok

On the Baire Category Theorem[1]

Artur Korniłowicz
Warsaw University
Białystok

Summary. In this paper Exercise 3.43 from Chapter 1 of [14] is solved.

MML Identifier: `WAYBEL12`.

WWW: `http://mizar.org/JFM/Vol9/waybel12.html`

The articles [21], [11], [25], [23], [26], [9], [10], [7], [13], [8], [1], [2], [19], [24], [27], [12], [16], [20], [3], [4], [15], [5], [28], [17], [6], [18], and [22] provide the notation and terminology for this paper.

1. Preliminaries

Let T be a topological structure and let P be a subset of T. Let us observe that P is closed if and only if:

(Def. 1) $P^{\rm c}$ is open.

Let T be a topological structure and let F be a family of subsets of T. We say that F is dense if and only if:

(Def. 2) For every subset X of T such that $X \in F$ holds X is dense.

Let us mention that there exists a 1-sorted structure which is empty.

Let S be an empty 1-sorted structure. One can check that the carrier of S is empty.

Let S be an empty 1-sorted structure. Note that every subset of S is empty.

Let us note that every set which is finite is also countable.

Let us mention that there exists a set which is empty.

Let S be a 1-sorted structure. One can verify that there exists a subset of S which is empty.

Let us note that there exists a set which is non empty and finite.

Let L be a non empty relational structure. Note that there exists a subset of L which is non empty and finite.

Let us observe that $\mathbb{N}$ is infinite.

One can verify that there exists a set which is infinite and countable.

Let S be a 1-sorted structure. One can verify that there exists a family of subsets of S which is empty.

One can prove the following propositions:

(2)[1] For all sets X, Y such that $\overline{\overline{X}} \leq \overline{\overline{Y}}$ and Y is countable holds X is countable.

[1]This work has been partially supported by the Office of Naval Research Grant N00014-95-1-1336.

[1] The proposition (1) has been removed.

(3) For every infinite countable set A holds $\mathbb{N} \approx A$.

(4) For every non empty countable set A there exists a function f from $\mathbb{N}$ into A such that $\operatorname{rng} f = A$.

(5) For every 1-sorted structure S and for all subsets X, Y of S holds $(X \cup Y)^{\rm c} = X^{\rm c} \cap Y^{\rm c}$.

(6) For every 1-sorted structure S and for all subsets X, Y of S holds $(X \cap Y)^{\rm c} = X^{\rm c} \cup Y^{\rm c}$.

(7) Let L be a non empty transitive relational structure and A, B be subsets of L. If A is finer than B, then $\downarrow A \subseteq \downarrow B$.

(8) Let L be a non empty transitive relational structure and A, B be subsets of L. If A is coarser than B, then $\uparrow A \subseteq \uparrow B$.

(9) Let L be a non empty poset and D be a non empty finite filtered subset of L. If $\inf D$ exists in L, then $\inf D \in D$.

(10) Let L be a lower-bounded antisymmetric non empty relational structure and X be a non empty lower subset of L. Then $\perp_L \in X$.

(11) Let L be a lower-bounded antisymmetric non empty relational structure and X be a non empty subset of L. Then $\perp_L \in \downarrow X$.

(12) Let L be an upper-bounded antisymmetric non empty relational structure and X be a non empty upper subset of L. Then $\top_L \in X$.

(13) Let L be an upper-bounded antisymmetric non empty relational structure and X be a non empty subset of L. Then $\top_L \in \uparrow X$.

(14) Let L be a lower-bounded antisymmetric relational structure with g.l.b.'s and X be a subset of L. Then $X \sqcap \{\perp_L\} \subseteq \{\perp_L\}$.

(15) Let L be a lower-bounded antisymmetric relational structure with g.l.b.'s and X be a non empty subset of L. Then $X \sqcap \{\perp_L\} = \{\perp_L\}$.

(16) Let L be an upper-bounded antisymmetric relational structure with l.u.b.'s and X be a subset of L. Then $X \sqcup \{\top_L\} \subseteq \{\top_L\}$.

(17) Let L be an upper-bounded antisymmetric relational structure with l.u.b.'s and X be a non empty subset of L. Then $X \sqcup \{\top_L\} = \{\top_L\}$.

(18) For every upper-bounded semilattice L and for every subset X of L holds $\{\top_L\} \sqcap X = X$.

(19) For every lower-bounded poset L with l.u.b.'s and for every subset X of L holds $\{\perp_L\} \sqcup X = X$.

(20) Let L be a non empty reflexive relational structure and A, B be subsets of L. If $A \subseteq B$, then A is finer than B and coarser than B.

(21) Let L be an antisymmetric transitive relational structure with g.l.b.'s, V be a subset of L, and x, y be elements of L. If $x \leq y$, then $\{y\} \sqcap V$ is coarser than $\{x\} \sqcap V$.

(22) Let L be an antisymmetric transitive relational structure with l.u.b.'s, V be a subset of L, and x, y be elements of L. If $x \leq y$, then $\{x\} \sqcup V$ is finer than $\{y\} \sqcup V$.

(23) Let L be a non empty relational structure and V, S, T be subsets of L. If S is coarser than T and V is upper and $T \subseteq V$, then $S \subseteq V$.

(24) Let L be a non empty relational structure and V, S, T be subsets of L. If S is finer than T and V is lower and $T \subseteq V$, then $S \subseteq V$.

(25) For every semilattice L and for every upper filtered subset F of L holds $F \sqcap F = F$.

(26) For every sup-semilattice L and for every lower directed subset I of L holds $I \sqcup I = I$.

(27) For every upper-bounded semilattice L and for every subset V of L holds $\{x; x$ ranges over elements of L: $V \sqcap \{x\} \subseteq V\}$ is non empty.

(28) Let L be an antisymmetric transitive relational structure with g.l.b.'s and V be a subset of L. Then $\{x; x$ ranges over elements of L: $V \sqcap \{x\} \subseteq V\}$ is a filtered subset of L.

(29) Let L be an antisymmetric transitive relational structure with g.l.b.'s and V be an upper subset of L. Then $\{x; x$ ranges over elements of L: $V \sqcap \{x\} \subseteq V\}$ is an upper subset of L.

(30) For every poset L with g.l.b.'s and for every subset X of L such that X is open and lower holds X is filtered.

Let L be a poset with g.l.b.'s. Observe that every subset of L which is open and lower is also filtered.

Let L be a continuous antisymmetric non empty reflexive relational structure. Note that every subset of L which is lower is also open.

Let L be a continuous semilattice and let x be an element of L. Note that $(\downarrow x)^{\mathrm{c}}$ is open.

One can prove the following two propositions:

(31) Let L be a semilattice and C be a non empty subset of L. Suppose that for all elements x, y of L such that $x \in C$ and $y \in C$ holds $x \leq y$ or $y \leq x$. Let Y be a non empty finite subset of C. Then $\bigsqcap_L Y \in Y$.

(32) Let L be a sup-semilattice and C be a non empty subset of L. Suppose that for all elements x, y of L such that $x \in C$ and $y \in C$ holds $x \leq y$ or $y \leq x$. Let Y be a non empty finite subset of C. Then $\bigsqcup_L Y \in Y$.

Let L be a semilattice and let F be a filter of L. A subset of L is called a generator set of F if:

(Def. 3) $F = \uparrow \mathrm{fininfs}(\mathrm{it})$.

Let L be a semilattice and let F be a filter of L. Observe that there exists a generator set of F which is non empty.

We now state four propositions:

(33) Let L be a semilattice, A be a subset of L, and B be a non empty subset of L. If A is coarser than B, then $\mathrm{fininfs}(A)$ is coarser than $\mathrm{fininfs}(B)$.

(34) Let L be a semilattice, F be a filter of L, G be a generator set of F, and A be a non empty subset of L. Suppose G is coarser than A and A is coarser than F. Then A is a generator set of F.

(35) Let L be a semilattice, A be a subset of L, and f, g be functions from $\mathbb{N}$ into the carrier of L. Suppose $\mathrm{rng}\, f = A$ and for every element n of $\mathbb{N}$ holds $g(n) = \bigsqcap_L \{f(m); m$ ranges over natural numbers: $m \leq n\}$. Then A is coarser than $\mathrm{rng}\, g$.

(36) Let L be a semilattice, F be a filter of L, G be a generator set of F, and f, g be functions from $\mathbb{N}$ into the carrier of L. Suppose $\mathrm{rng}\, f = G$ and for every element n of $\mathbb{N}$ holds $g(n) = \bigsqcap_L \{f(m); m$ ranges over natural numbers: $m \leq n\}$. Then $\mathrm{rng}\, g$ is a generator set of F.

2. On the Baire Category Theorem

We now state four propositions:

(37) Let L be a lower-bounded continuous lattice, V be an open upper subset of L, F be a filter of L, and v be an element of L. Suppose $V \sqcap F \subseteq V$ and $v \in V$ and there exists a non empty generator set of F which is countable. Then there exists an open filter O of L such that $O \subseteq V$ and $v \in O$ and $F \subseteq O$.

(38) Let L be a lower-bounded continuous lattice, V be an open upper subset of L, N be a non empty countable subset of L, and v be an element of L. Suppose $V \sqcap N \subseteq V$ and $v \in V$. Then there exists an open filter O of L such that $\{v\} \sqcap N \subseteq O$ and $O \subseteq V$ and $v \in O$.

(39) Let L be a lower-bounded continuous lattice, V be an open upper subset of L, N be a non empty countable subset of L, and x, y be elements of L. Suppose $V \sqcap N \subseteq V$ and $y \in V$ and $x \notin V$. Then there exists an irreducible element p of L such that $x \leq p$ and $p \notin \uparrow(\{y\} \sqcap N)$.

(40) Let L be a lower-bounded continuous lattice, x be an element of L, and N be a non empty countable subset of L. Suppose that for all elements n, y of L such that $y \not\leq x$ and $n \in N$ holds $y \sqcap n \not\leq x$. Let y be an element of L. Suppose $y \not\leq x$. Then there exists an irreducible element p of L such that $x \leq p$ and $p \notin \uparrow(\{y\} \sqcap N)$.

Let L be a non empty relational structure and let u be an element of L. We say that u is dense if and only if:

(Def. 4) For every element v of L such that $v \neq \bot_L$ holds $u \sqcap v \neq \bot_L$.

Let L be an upper-bounded semilattice. Observe that $\top_L$ is dense.

Let L be an upper-bounded semilattice. Note that there exists an element of L which is dense.

The following proposition is true

(41) For every non trivial bounded semilattice L and for every element x of L such that x is dense holds $x \neq \bot_L$.

Let L be a non empty relational structure and let D be a subset of L. We say that D is dense if and only if:

(Def. 5) For every element d of L such that $d \in D$ holds d is dense.

The following proposition is true

(42) For every upper-bounded semilattice L holds $\{\top_L\}$ is dense.

Let L be an upper-bounded semilattice. Observe that there exists a subset of L which is non empty, finite, countable, and dense.

Next we state several propositions:

(43) Let L be a lower-bounded continuous lattice, D be a non empty countable dense subset of L, and u be an element of L. Suppose $u \neq \bot_L$. Then there exists an irreducible element p of L such that $p \neq \top_L$ and $p \notin \uparrow(\{u\} \sqcap D)$.

(44) Let T be a non empty topological space, A be an element of $\langle$the topology of T, $\subseteq\rangle$, and B be a subset of T. If $A = B$ and B^{c} is irreducible, then A is irreducible.

(45) Let T be a non empty topological space, A be an element of $\langle$the topology of T, $\subseteq\rangle$, and B be a subset of T. Suppose $A = B$ and $A \neq \top_{\langle \text{the topology of } T, \subseteq \rangle}$. Then A is irreducible if and only if B^{c} is irreducible.

(46) Let T be a non empty topological space, A be an element of $\langle$the topology of T, $\subseteq\rangle$, and B be a subset of T. If $A = B$, then A is dense iff B is everywhere dense.

(47) Let T be a non empty topological space. Suppose T is locally-compact. Let D be a countable family of subsets of T. Suppose D is non empty, dense, and open. Let O be a non empty subset of T. Suppose O is open. Then there exists an irreducible subset A of T such that for every subset V of T if $V \in D$, then $A \cap O$ meets V.

Let T be a non empty topological space. Let us observe that T is Baire if and only if the condition (Def. 6) is satisfied.

(Def. 6) Let F be a family of subsets of T. Suppose F is countable and for every subset S of T such that $S \in F$ holds S is open and dense. Then there exists a subset I of T such that $I = \mathrm{Intersect}(F)$ and I is dense.

Next we state the proposition

(48) For every non empty topological space T such that T is sober and locally-compact holds T is Baire.

REFERENCES

[1] Grzegorz Bancerek. Cardinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/card_1.html.

[2] Grzegorz Bancerek. Countable sets and Hessenberg's theorem. *Journal of Formalized Mathematics*, 2, 1990. http://mizar.org/JFM/Vol2/card_4.html.

[3] Grzegorz Bancerek. Complete lattices. *Journal of Formalized Mathematics*, 4, 1992. http://mizar.org/JFM/Vol4/lattice3.html.

[4] Grzegorz Bancerek. Bounds in posets and relational substructures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_0.html.

[5] Grzegorz Bancerek. Directed sets, nets, ideals, filters, and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_0.html.

[6] Grzegorz Bancerek. The "way-below" relation. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_3.html.

[7] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finseq_1.html.

[8] Józef Białas. Group and field definitions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/realset1.html.

[9] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[10] Czesław Byliński. Functions from a set to a set. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_2.html.

[11] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[12] Agata Darmochwał. Families of subsets, subspaces and mappings in topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_2.html.

[13] Agata Darmochwał. Finite sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/finset_1.html.

[14] G. Gierz, K.H. Hofmann, K. Keimel, J.D. Lawson, M. Mislove, and D.S. Scott. *A Compendium of Continuous Lattices*. Springer-Verlag, Berlin, Heidelberg, New York, 1980.

[15] Adam Grabowski and Robert Milewski. Boolean posets, posets under inclusion and products of relational structures. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_1.html.

[16] Zbigniew Karno. Remarks on special subsets of topological spaces. *Journal of Formalized Mathematics*, 5, 1993. http://mizar.org/JFM/Vol5/tops_3.html.

[17] Artur Korniłowicz. Definitions and properties of the join and meet of subsets. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_4.html.

[18] Beata Madras. Irreducible and prime elements. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/waybel_6.html.

[19] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/pre_topc.html.

[20] Alexander Yu. Shibakov and Andrzej Trybulec. The Cantor set. *Journal of Formalized Mathematics*, 7, 1995. http://mizar.org/JFM/Vol7/cantor_1.html.

[21] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[22] Andrzej Trybulec. Baire spaces, Sober spaces. *Journal of Formalized Mathematics*, 9, 1997. http://mizar.org/JFM/Vol9/yellow_8.html.

[23] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[24] Wojciech A. Trybulec. Partially ordered sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/orders_1.html.

[25] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

[26] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

[27] Mirosław Wysocki and Agata Darmochwał. Subsets of topological spaces. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/tops_1.html.

[28] Mariusz Żynel and Czesław Byliński. Properties of relational structures, posets, lattices and maps. *Journal of Formalized Mathematics*, 8, 1996. http://mizar.org/JFM/Vol8/yellow_2.html.

Received February 5, 1997

Published January 2, 2004