JOURNAL OF FORMALIZED MATHEMATICS
Volume 1, Released 1989, Published 2003
Inst. of Computer Science, Univ. of Białystok

Zermelo Theorem and Axiom of Choice

Grzegorz Bancerek
Warsaw University
Białystok

Summary. The article is continuation of [2] and [1], and the goal of it is show that Zermelo theorem (every set has a relation which well orders it - proposition (26)) and axiom of choice (for every non-empty family of non-empty and separate sets there is set which has exactly one common element with arbitrary family member - proposition (27)) are true. It is result of the Tarski's axiom A introduced in [5] and repeated in [6]. Inclusion as a settheoretical binary relation is introduced, the correspondence of well ordering relations to ordinal numbers is shown, and basic properties of equinumerosity are presented. Some facts are based on [4].

MML Identifier: WELLORD2.

WWW: http://mizar.org/JFM/Vol1/wellord2.html

The articles [6], [7], [8], [3], [2], and [1] provide the notation and terminology for this paper.

For simplicity, we adopt the following rules: X, Y, Z denote sets, a, x denote sets, R denotes a binary relation, and A, B denote ordinal numbers.

Let us consider X. The functor ${}^{\subseteq}_{X}$ yielding a binary relation is defined as follows:

(Def. 1) $\operatorname{field}({}^{\subseteq}_{X}) = X$ and for all Y, Z such that $Y \in X$ and $Z \in X$ holds $\langle Y, Z\rangle \in {}^{\subseteq}_{X}$ iff $Y \subseteq Z$.

One can prove the following propositions:

(2)[1] ${}^{\subseteq}_{X}$ is reflexive.

(3) ${}^{\subseteq}_{X}$ is transitive.

(4) ${}^{\subseteq}_{A}$ is connected.

(5) ${}^{\subseteq}_{X}$ is antisymmetric.

(6) ${}^{\subseteq}_{A}$ is well founded.

(7) ${}^{\subseteq}_{A}$ is well-ordering.

(8) If $Y \subseteq X$, then ${}^{\subseteq}_{X} \upharpoonright^2 Y = {}^{\subseteq}_{Y}$.

(9) For all A, X such that $X \subseteq A$ holds ${}^{\subseteq}_{X}$ is well-ordering.

(10) If $A \in B$, then $A = {}^{\subseteq}_{B}\text{-Seg}(A)$.

(11) If ${}^{\subseteq}_{A}$ and ${}^{\subseteq}_{B}$ are isomorphic, then $A = B$.

(12) For all X, R, A, B such that R and ${}^{\subseteq}_{A}$ are isomorphic and R and ${}^{\subseteq}_{B}$ are isomorphic holds $A = B$.

[1] The proposition (1) has been removed.

© Association of Mizar Users

(13) Let given R. Suppose R is well-ordering and for every a such that $a \in \operatorname{field} R$ there exists A such that $R \mid^2 R\text{-Seg}(a)$ and $\subseteq_A$ are isomorphic. Then there exists A such that R and $\subseteq_A$ are isomorphic.

(14) For every R such that R is well-ordering there exists A such that R and $\subseteq_A$ are isomorphic.

Let us consider R. Let us assume that R is well-ordering. The functor $\overline{R}$ yields an ordinal number and is defined as follows:

(Def. 2) R and $\subseteq_{\overline{R}}$ are isomorphic.

Let us consider A, R. We say that A is an order type of R if and only if:

(Def. 3) $A = \overline{R}$.

One can prove the following proposition

(17)[2] If $X \subseteq A$, then $\overline{\subseteq_X} \subseteq A$.

In the sequel f denotes a function.

Let us consider X, Y. Let us observe that $X \approx Y$ if and only if:

(Def. 4) There exists f such that f is one-to-one and $\operatorname{dom} f = X$ and $\operatorname{rng} f = Y$.

Let us notice that the predicate $X \approx Y$ is reflexive and symmetric.

We now state three propositions:

(22)[3] If $X \approx Y$ and $Y \approx Z$, then $X \approx Z$.

(25)[4] If R well orders X, then $\operatorname{field}(R \mid^2 X) = X$ and $R \mid^2 X$ is well-ordering.

(26) For every X there exists R such that R well orders X.

In the sequel M denotes a non empty set.

Next we state two propositions:

(27) Suppose for every X such that $X \in M$ holds $X \neq \emptyset$ and for all X, Y such that $X \in M$ and $Y \in M$ and $X \neq Y$ holds X misses Y. Then there exists a set C_1 such that for every X such that $X \in M$ there exists x such that $C_1 \cap X = \{x\}$.

(28) If for every X such that $X \in M$ holds $X \neq \emptyset$, then there exists a function C_1 such that $\operatorname{dom} C_1 = M$ and for every X such that $X \in M$ holds $C_1(X) \in X$.

REFERENCES

[1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[2] Grzegorz Bancerek. The well ordering relations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/wellord1.html.

[3] Czesław Byliński. Functions and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/funct_1.html.

[4] Kazimierz Kuratowski and Andrzej Mostowski. *Teoria mnogości*. PTM, Wrocław, 1952.

[5] Alfred Tarski. Über Unerreichbare Kardinalzahlen. *Fundamenta Mathematicae*, 30:176–183, 1938.

[6] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[7] Edmund Woronowicz. Relations and their basic properties. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_1.html.

[2] The propositions (15) and (16) have been removed.

[3] The propositions (18)–(21) have been removed.

[4] The propositions (23) and (24) have been removed.

[8] Edmund Woronowicz and Anna Zalewska. Properties of binary relations. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/relat_2.html.

Received June 26, 1989

Published January 2, 2004