JOURNAL OF FORMALIZED MATHEMATICS
Volume EMM, Released 2003, Published 2003
Inst. of Computer Science, Univ. of Białystok

Introduction to Arithmetic of Real Numbers

Library Committee
Association of Mizar Users

MML Identifier: XREAL_0.

WWW: http://mizar.org/JFM/EMM/xreal_0.html

The articles [4], [3], [7], [1], [2], [5], and [6] provide the notation and terminology for this paper.

1. Main Definitions

Let r be a number. We say that r is real if and only if:

(Def. 1) $r \in \mathbb{R}$.

Let us mention that every number which is natural is also real and every number which is real is also complex.

Let us note that there exists a number which is real.

Let x, y be real numbers. The predicate $x \leq y$ is defined by:

(Def. 2)(i) There exist elements x', y' of $\mathbb{R}_+$ such that $x = x'$ and $y = y'$ and $x' \leq y'$ if $x \in \mathbb{R}_+$ and $y \in \mathbb{R}_+$,

(ii) there exist elements x', y' of $\mathbb{R}_+$ such that $x = \langle 0, x' \rangle$ and $y = \langle 0, y' \rangle$ and $y' \leq x'$ if $x \in [:\{0\}, \mathbb{R}_+:]$ and $y \in [:\{0\}, \mathbb{R}_+:]$,

(iii) $y \in \mathbb{R}_+$ and $x \in [:\{0\}, \mathbb{R}_+:]$, otherwise.

Let us notice that the predicate $x \leq y$ is reflexive and connected. We introduce $y \geq x$ as a synonym of $x \leq y$. We introduce $y < x$ and $x > y$ as antonyms of $x \leq y$.

Let x be a real number. We say that x is positive if and only if:

(Def. 3) $x > 0$.

We say that x is negative if and only if:

(Def. 4) $x < 0$.

Let x be a real number. One can verify that $-x$ is real and x^{-1} is real. Let y be a real number. Observe that $x + y$ is real and $x \cdot y$ is real.

Let x, y be real numbers. Note that $x - y$ is real and $\frac{x}{y}$ is real.

2. Clusters

One can check the following observations:

* every real number which is positive is also non negative and non zero,
* every real number which is non negative and non zero is also positive,

© Association of Mizar Users

* every real number which is negative is also non positive and non zero,
* every real number which is non positive and non zero is also negative,
* every real number which is zero is also non negative and non positive, and
* every real number which is non negative and non positive is also zero.

One can check the following observations:

* there exists a real number which is positive,
* there exists a real number which is negative, and
* there exists a real number which is zero.

Let r, s be non negative real numbers. Note that $r+s$ is non negative.

Let r, s be non positive real numbers. Note that $r+s$ is non positive.

Let r be a positive real number and let s be a non negative real number. One can verify that $r+s$ is positive and $s+r$ is positive.

Let r be a negative real number and let s be a non positive real number. Note that $r+s$ is negative and $s+r$ is negative.

Let r be a non positive real number. Observe that $-r$ is non negative.

Let r be a non negative real number. One can verify that $-r$ is non positive.

Let r be a non negative real number and let s be a non positive real number. One can check that $r-s$ is non negative and $s-r$ is non positive.

Let r be a positive real number and let s be a non positive real number. Note that $r-s$ is positive and $s-r$ is negative.

Let r be a negative real number and let s be a non negative real number. Note that $r-s$ is negative and $s-r$ is positive.

Let r be a non positive real number and let s be a non negative real number. Observe that $r \cdot s$ is non positive and $s \cdot r$ is non positive.

Let r, s be non positive real numbers. One can check that $r \cdot s$ is non negative.

Let r, s be non negative real numbers. Note that $r \cdot s$ is non negative.

Let r be a non positive real number. Note that r^{-1} is non positive.

Let r be a non negative real number. One can check that r^{-1} is non negative.

Let r be a non negative real number and let s be a non positive real number. Note that $\frac{r}{s}$ is non positive and $\frac{s}{r}$ is non positive.

Let r, s be non negative real numbers. One can verify that $\frac{r}{s}$ is non negative.

Let r, s be non positive real numbers. Note that $\frac{r}{s}$ is non negative.

REFERENCES

[1] Grzegorz Bancerek. The ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal1.html.

[2] Grzegorz Bancerek. Sequences of ordinal numbers. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/ordinal2.html.

[3] Czesław Byliński. Some basic properties of sets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/zfmisc_1.html.

[4] Andrzej Trybulec. Tarski Grothendieck set theory. *Journal of Formalized Mathematics*, Axiomatics, 1989. http://mizar.org/JFM/Axiomatics/tarski.html.

[5] Andrzej Trybulec. Non negative real numbers. Part I. *Journal of Formalized Mathematics*, Addenda, 1998. http://mizar.org/JFM/Addenda/arytm_2.html.

[6] Andrzej Trybulec. Subsets of real numbers. *Journal of Formalized Mathematics*, Addenda, 2003. http://mizar.org/JFM/Addenda/numbers.html.

[7] Zinaida Trybulec. Properties of subsets. *Journal of Formalized Mathematics*, 1, 1989. http://mizar.org/JFM/Vol1/subset_1.html.

Received February 11, 2003

Published January 2, 2004