

Boolean Properties of Lattices

Agnieszka Julia Marasik
Warsaw University
Białystok

MML Identifier: BOOLEALG.

The article [1] provides the terminology and notation for this paper.

1. GENERAL LATTICE

We follow the rules: L will be a lattice and X, Y, Z, V will be elements of the carrier of L .

Let us consider L, X, Y . The functor $X \setminus Y$ yielding an element of the carrier of L is defined by:

$$(Def.1) \quad X \setminus Y = X \sqcap Y^c.$$

Let us consider L, X, Y . The functor $X \dot{-} Y$ yields an element of the carrier of L and is defined by:

$$(Def.2) \quad X \dot{-} Y = (X \setminus Y) \sqcup (Y \setminus X).$$

Let us consider L, X, Y . Let us observe that $X = Y$ if and only if:

$$(Def.3) \quad X \sqsubseteq Y \text{ and } Y \sqsubseteq X.$$

Let us consider L, X, Y . We say that X meets Y if and only if:

$$(Def.4) \quad X \sqcap Y \neq \perp_L.$$

We introduce X misses Y as an antonym of X meets Y .

We now state a number of propositions:

- (1) $X \sqsubseteq X \sqcup Y$ and $Y \sqsubseteq X \sqcup Y$.
- (3)¹ If $X \sqcup Y \sqsubseteq Z$, then $X \sqsubseteq Z$ and $Y \sqsubseteq Z$.
- (4) $X \sqcap Y \sqsubseteq X \sqcup Z$.
- (5) If $X \sqsubseteq Y$, then $X \sqcap Z \sqsubseteq Y \sqcap Z$ and $Z \sqcap X \sqsubseteq Z \sqcap Y$.
- (6) If $X \sqsubseteq Z$, then $X \setminus Y \sqsubseteq Z$.

¹The proposition (2) has been removed.

- (7) If $X \sqsubseteq Y$, then $X \setminus Z \sqsubseteq Y \setminus Z$.
- (8) $X \setminus Y \sqsubseteq X$.
- (9) $X \setminus Y \sqsubseteq X \dot{-} Y$.
- (10) If $X \setminus Y \sqsubseteq Z$ and $Y \setminus X \sqsubseteq Z$, then $X \dot{-} Y \sqsubseteq Z$.
- (11) $X = Y \sqcup Z$ iff $Y \sqsubseteq X$ and $Z \sqsubseteq X$ and for every V such that $Y \sqsubseteq V$ and $Z \sqsubseteq V$ holds $X \sqsubseteq V$.
- (12) $X = Y \sqcap Z$ iff $X \sqsubseteq Y$ and $X \sqsubseteq Z$ and for every V such that $V \sqsubseteq Y$ and $V \sqsubseteq Z$ holds $V \sqsubseteq X$.
- (13) If $X \sqcup Y = Y$ or $Y \sqcup X = Y$, then $X \sqsubseteq Y$.
- (14) $X \sqcap (Y \setminus Z) = X \sqcap Y \setminus Z$.
- (15) If X meets Y , then Y meets X .
- (16) X meets X iff $X \neq \perp_L$.
- (17) $X \dot{-} Y = Y \dot{-} X$.

2. MODULAR LATTICE

In the sequel L will denote a modular lattice and X, Y will denote elements of the carrier of L .

The following three propositions are true:

- (18) If $Y \sqsubseteq X$ and $X \sqcap Y = \perp_L$, then $Y = \perp_L$.
- (20)² If $X \sqsubseteq Y$, then $X \sqcup Y = Y$ and $Y \sqcup X = Y$.
- (21) If X misses Y , then Y misses X .

3. DISTRIBUTIVE LATTICE

In the sequel L will denote a distributive lattice and X, Y, Z will denote elements of the carrier of L .

Next we state three propositions:

- (22) If $X \sqcap Y \sqcup X \sqcap Z = X$, then $X \sqsubseteq Y \sqcup Z$.
- (23) $X \sqcap Y \sqcup Y \sqcap Z \sqcup Z \sqcap X = (X \sqcup Y) \sqcap (Y \sqcup Z) \sqcap (Z \sqcup X)$.
- (24) $(X \sqcup Y) \setminus Z = (X \setminus Z) \sqcup (Y \setminus Z)$.

²The proposition (19) has been removed.

4. DISTRIBUTIVE LOWER BOUNDED LATTICE

In the sequel L will denote a lower bound lattice and X, Y, Z will denote elements of the carrier of L .

The following propositions are true:

- (25) If $X \sqsubseteq \perp_L$, then $X = \perp_L$.
- (26) If $X \sqsubseteq Y$ and $X \sqsubseteq Z$ and $Y \sqcap Z = \perp_L$, then $X = \perp_L$.
- (27) $X \sqcup Y = \perp_L$ iff $X = \perp_L$ and $Y = \perp_L$.
- (28) If $X \sqsubseteq Y$ and $Y \sqcap Z = \perp_L$, then $X \sqcap Z = \perp_L$.
- (29) $\perp_L \setminus X = \perp_L$.
- (30) If X meets Y and $Y \sqsubseteq Z$, then X meets Z .
- (31) If X meets $Y \sqcap Z$, then X meets Y and X meets Z .
- (32) If X meets $Y \setminus Z$, then X meets Y .
- (33) X misses \perp_L .
- (34) If X misses Z and $Y \sqsubseteq Z$, then X misses Y .
- (35) If X misses Y or X misses Z , then X misses $Y \sqcap Z$.
- (36) If $X \sqsubseteq Y$ and $X \sqsubseteq Z$ and Y misses Z , then $X = \perp_L$.
- (37) If X misses Y , then $Z \sqcap X$ misses $Z \sqcap Y$ and $X \sqcap Z$ misses $Y \sqcap Z$.

5. BOOLEAN LATTICE

We follow a convention: L will be a Boolean lattice and X, Y, Z, V will be elements of the carrier of L .

Next we state a number of propositions:

- (38) If $X \setminus Y \sqsubseteq Z$, then $X \sqsubseteq Y \sqcup Z$.
- (39) If $X \sqsubseteq Y$, then $Z \setminus Y \sqsubseteq Z \setminus X$.
- (40) If $X \sqsubseteq Y$ and $Z \sqsubseteq V$, then $X \setminus V \sqsubseteq Y \setminus Z$.
- (41) If $X \sqsubseteq Y \sqcup Z$, then $X \setminus Y \sqsubseteq Z$ and $X \setminus Z \sqsubseteq Y$.
- (42) $X^c \sqsubseteq (X \sqcap Y)^c$ and $Y^c \sqsubseteq (X \sqcap Y)^c$.
- (43) $(X \sqcup Y)^c \sqsubseteq X^c$ and $(X \sqcup Y)^c \sqsubseteq Y^c$.
- (44) If $X \sqsubseteq Y \setminus X$, then $X = \perp_L$.
- (45) If $X \sqsubseteq Y$, then $Y = X \sqcup (Y \setminus X)$ and $Y = (Y \setminus X) \sqcup X$.
- (46) $X \setminus Y = \perp_L$ iff $X \sqsubseteq Y$.
- (47) If $X \sqsubseteq Y \sqcup Z$ and $X \sqcap Z = \perp_L$, then $X \sqsubseteq Y$.
- (48) $X \sqcup Y = (X \setminus Y) \sqcup Y$.
- (49) $X \setminus (X \sqcup Y) = \perp_L$ and $X \setminus (Y \sqcup X) = \perp_L$.
- (50) $X \setminus X \sqcap Y = X \setminus Y$ and $X \setminus Y \sqcap X = X \setminus Y$.
- (51) $(X \setminus Y) \sqcap Y = \perp_L$ and $Y \sqcap (X \setminus Y) = \perp_L$.

- (52) $X \sqcup (Y \setminus X) = X \sqcup Y$ and $(Y \setminus X) \sqcup X = Y \sqcup X$.
- (53) $X \sqcap Y \sqcup (X \setminus Y) = X$ and $(X \setminus Y) \sqcup X \sqcap Y = X$.
- (54) $X \setminus (Y \setminus Z) = (X \setminus Y) \sqcup X \sqcap Z$.
- (55) $X \setminus (X \setminus Y) = X \sqcap Y$.
- (56) $(X \sqcup Y) \setminus Y = X \setminus Y$.
- (57) $X \sqcap Y = \perp_L$ iff $X \setminus Y = X$.
- (58) $X \setminus (Y \sqcup Z) = (X \setminus Y) \sqcap (X \setminus Z)$.
- (59) $X \setminus Y \sqcap Z = (X \setminus Y) \sqcup (X \setminus Z)$.
- (60) $X \sqcap (Y \setminus Z) = X \sqcap Y \setminus X \sqcap Z$ and $(Y \setminus Z) \sqcap X = Y \sqcap X \setminus Z \sqcap X$.
- (61) $(X \sqcup Y) \setminus X \sqcap Y = (X \setminus Y) \sqcup (Y \setminus X)$.
- (62) $X \setminus Y \setminus Z = X \setminus (Y \sqcup Z)$.
- (63) If $X \setminus Y = Y \setminus X$, then $X = Y$.
- (64) $(\perp_L)^c = \top_L$.
- (65) $(\top_L)^c = \perp_L$.
- (66) $X \setminus X = \perp_L$.
- (67) $X \setminus \perp_L = X$.
- (68) $(X \setminus Y)^c = X^c \sqcup Y$.
- (69) X meets $Y \sqcup Z$ iff X meets Y or X meets Z .
- (70) $X \sqcap Y$ misses $X \setminus Y$.
- (71) X misses $Y \sqcup Z$ iff X misses Y and X misses Z .
- (72) $X \setminus Y$ misses Y .
- (73) If X misses Y , then $(X \sqcup Y) \setminus Y = X$ and $(X \sqcup Y) \setminus X = Y$.
- (74) If $X^c \sqcup Y^c = X \sqcup Y$ and X misses X^c and Y misses Y^c , then $X = Y^c$ and $Y = X^c$.
- (75) If $X^c \sqcup Y^c = X \sqcup Y$ and Y misses X^c and X misses Y^c , then $X = X^c$ and $Y = Y^c$.
- (76) $X \dot{\sqcup} \perp_L = X$ and $\perp_L \dot{\sqcup} X = X$.
- (77) $X \dot{\sqcup} X = \perp_L$.
- (78) $X \sqcap Y$ misses $X \dot{\sqcup} Y$.
- (79) $X \sqcup Y = X \dot{\sqcup} (Y \setminus X)$.
- (80) $X \dot{\sqcup} X \sqcap Y = X \setminus Y$.
- (81) $X \sqcup Y = (X \dot{\sqcup} Y) \sqcup X \sqcap Y$.
- (82) $X \dot{\sqcup} Y \dot{\sqcup} X \sqcap Y = X \sqcup Y$.
- (83) $X \dot{\sqcup} Y \dot{\sqcup} (X \sqcup Y) = X \sqcap Y$.
- (84) $X \dot{\sqcup} Y = (X \sqcup Y) \setminus X \sqcap Y$.
- (85) $(X \dot{\sqcup} Y) \setminus Z = (X \setminus (Y \sqcup Z)) \sqcup (Y \setminus (X \sqcup Z))$.
- (86) $X \setminus (Y \dot{\sqcup} Z) = (X \setminus (Y \sqcup Z)) \sqcup X \sqcap Y \sqcap Z$.
- (87) $(X \dot{\sqcup} Y) \dot{\sqcup} Z = X \dot{\sqcup} (Y \dot{\sqcup} Z)$.
- (88) $(X \dot{\sqcup} Y)^c = X \sqcap Y \sqcup X^c \sqcap Y^c$.

REFERENCES

- [1] Stanisław Żukowski. Introduction to lattice theory. *Formalized Mathematics*, 1(1):215–222, 1990.

Received March 28, 1994
