

# Higher-Order Partial Differentiation<sup>1</sup>

Noboru Endou  
 Nagano National College of Technology  
 Japan

Hiroyuki Okazaki  
 Shinshu University  
 Nagano, Japan

Yasunari Shidama  
 Shinshu University  
 Nagano, Japan

**Summary.** In this article, we shall extend the formalization of [10] to discuss higher-order partial differentiation of real valued functions. The linearity of this operator is also proved (refer to [10], [12] and [13] for partial differentiation).

MML identifier: PDIFF\_9, version: 7.12.02 4.181.1147

The terminology and notation used here have been introduced in the following articles: [3], [8], [2], [4], [5], [15], [21], [17], [16], [20], [1], [6], [10], [12], [13], [18], [11], [9], [23], [7], [19], [14], and [22].

## 1. PRELIMINARIES

We use the following convention:  $m, n$  denote non empty elements of  $\mathbb{N}$ ,  $i, j$  denote elements of  $\mathbb{N}$ , and  $Z$  denotes a set.

One can prove the following propositions:

- (1) Let  $S, T$  be real normed spaces,  $f$  be a point of the real norm space of bounded linear operators from  $S$  into  $T$ , and  $r$  be a real number. Suppose  $0 \leq r$  and for every point  $x$  of  $S$  such that  $\|x\| \leq 1$  holds  $\|f(x)\| \leq r \cdot \|x\|$ . Then  $\|f\| \leq r$ .
- (2) Let  $S$  be a real normed space and  $f$  be a partial function from  $S$  to  $\mathbb{R}$ . Then  $f$  is continuous on  $Z$  if and only if the following conditions are satisfied:

---

<sup>1</sup>This work was supported by JSPS KAKENHI 22300285 and 23500029.

- (i)  $Z \subseteq \text{dom } f$ , and
- (ii) for every sequence  $s_1$  of  $S$  such that  $\text{rng } s_1 \subseteq Z$  and  $s_1$  is convergent and  $\lim s_1 \in Z$  holds  $f_* s_1$  is convergent and  $f_{\lim s_1} = \lim(f_* s_1)$ .
- (3) For every partial function  $f$  from  $\mathcal{R}^i$  to  $\mathbb{R}$  holds  $\text{dom} \langle f \rangle = \text{dom } f$ .
- (4) For every partial function  $f$  from  $\mathcal{R}^i$  to  $\mathbb{R}$  such that  $Z \subseteq \text{dom } f$  holds  $\text{dom}(\langle f \rangle \upharpoonright Z) = Z$ .
- (5) For every partial function  $f$  from  $\mathcal{R}^i$  to  $\mathbb{R}$  holds  $\langle f \upharpoonright Z \rangle = \langle f \rangle \upharpoonright Z$ .
- (6) Let  $f$  be a partial function from  $\mathcal{R}^i$  to  $\mathbb{R}$  and  $x$  be an element of  $\mathcal{R}^i$ . If  $x \in \text{dom } f$ , then  $\langle f \rangle(x) = \langle f(x) \rangle$  and  $\langle f \rangle_x = \langle f_x \rangle$ .
- (7) For all partial functions  $f, g$  from  $\mathcal{R}^i$  to  $\mathbb{R}$  holds  $\langle f + g \rangle = \langle f \rangle + \langle g \rangle$  and  $\langle f - g \rangle = \langle f \rangle - \langle g \rangle$ .
- (8) For every partial function  $f$  from  $\mathcal{R}^i$  to  $\mathbb{R}$  and for every real number  $r$  holds  $\langle r \cdot f \rangle = r \cdot \langle f \rangle$ .
- (9) Let  $f$  be a partial function from  $\mathcal{R}^i$  to  $\mathbb{R}$  and  $g$  be a partial function from  $\mathcal{R}^i$  to  $\mathcal{R}^1$ . If  $\langle f \rangle = g$ , then  $|f| = |g|$ .
- (10) For every subset  $X$  of  $\mathcal{R}^m$  and for every subset  $Y$  of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  such that  $X = Y$  holds  $X$  is open iff  $Y$  is open.
- (11) For every element  $q$  of  $\mathbb{R}$  such that  $1 \leq i \leq j$  holds  $|\text{reproj}(i, \underbrace{\langle 0, \dots, 0 \rangle}_j)(q)| = |q|$ .
- (12) For every element  $x$  of  $\mathcal{R}^j$  holds  $x = (\text{reproj}(i, x))((\text{proj}(i, j))(x))$ .

## 2. CONTINUITY AND DIFFERENTIABILITY

The following two propositions are true:

- (13) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . If  $f$  is differentiable on  $X$ , then  $X$  is open.
- (14) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . Suppose  $X$  is open. Then  $f$  is differentiable on  $X$  if and only if the following conditions are satisfied:
  - (i)  $X \subseteq \text{dom } f$ , and
  - (ii) for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $f$  is differentiable in  $x$ .

Let  $m, n$  be non empty elements of  $\mathbb{N}$ , let  $Z$  be a set, and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . Let us assume that  $Z \subseteq \text{dom } f$ . The functor  $f'_{\upharpoonright Z}$  yields a partial function from  $\mathcal{R}^m$  to  $(\mathcal{R}^n)^{\mathcal{R}^m}$  and is defined by:

- (Def. 1)  $\text{dom}(f'_{\upharpoonright Z}) = Z$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in Z$  holds  $(f'_{\upharpoonright Z})_x = f'(x)$ .

We now state a number of propositions:

- (15) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . Suppose  $f$  is differentiable on  $X$  and  $g$  is differentiable on  $X$ . Then  $f + g$  is differentiable on  $X$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((f + g)'|_X)_x = f'(x) + g'(x)$ .
- (16) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . Suppose  $f$  is differentiable on  $X$  and  $g$  is differentiable on  $X$ . Then  $f - g$  is differentiable on  $X$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((f - g)'|_X)_x = f'(x) - g'(x)$ .
- (17) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ , and  $r$  be a real number. Suppose  $f$  is differentiable on  $X$ . Then  $r \cdot f$  is differentiable on  $X$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((r \cdot f)'|_X)_x = r \cdot f'(x)$ .
- (18) Let  $f$  be a point of the real norm space of bounded linear operators from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^j, \|\cdot\| \rangle$ . Then there exists a point  $p$  of  $\langle \mathcal{E}^j, \|\cdot\| \rangle$  such that
- (i)  $p = f(\langle 1 \rangle)$ ,
  - (ii) for every real number  $r$  and for every point  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  such that  $x = \langle r \rangle$  holds  $f(x) = r \cdot p$ , and
  - (iii) for every point  $x$  of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  holds  $\|f(x)\| = \|p\| \cdot \|x\|$ .
- (19) Let  $f$  be a point of the real norm space of bounded linear operators from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^j, \|\cdot\| \rangle$ . Then there exists a point  $p$  of  $\langle \mathcal{E}^j, \|\cdot\| \rangle$  such that  $p = f(\langle 1 \rangle)$  and  $\|p\| = \|f\|$ .
- (20) Let  $f$  be a point of the real norm space of bounded linear operators from  $\langle \mathcal{E}^1, \|\cdot\| \rangle$  into  $\langle \mathcal{E}^j, \|\cdot\| \rangle$  and  $x$  be a point of  $\langle \mathcal{E}^1, \|\cdot\| \rangle$ . Then  $\|f(x)\| = \|f\| \cdot \|x\|$ .
- (21) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $X$  be a subset of  $\mathcal{R}^m$ , and  $Y$  be a subset of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . Suppose  $1 \leq i \leq m$  and  $X$  is open and  $g = f$  and  $X = Y$  and  $f$  is partially differentiable on  $X$  w.r.t.  $i$ . Let  $x$  be an element of  $\mathcal{R}^m$  and  $y$  be a point of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . If  $x \in X$  and  $x = y$ , then  $\text{partdiff}(f, x, i) = (\text{partdiff}(g, y, i))(\langle 1 \rangle)$ .
- (22) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $X$  be a subset of  $\mathcal{R}^m$ , and  $Y$  be a subset of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . Suppose  $1 \leq i \leq m$  and  $X$  is open and  $g = f$  and  $X = Y$  and  $f$  is partially differentiable on  $X$  w.r.t.  $i$ . Let  $x_0, x_1$  be elements of  $\mathcal{R}^m$  and  $y_0, y_1$  be points of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . If  $x_0 = y_0$  and  $x_1 = y_1$  and  $x_0, x_1 \in X$ , then  $\|(f|_X^i)_{x_1} - (f|_X^i)_{x_0}\| = \|(g|_Y^i)_{y_1} - (g|_Y^i)_{y_0}\|$ .
- (23) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $X$  be a subset of  $\mathcal{R}^m$ , and  $Y$  be a subset of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . Suppose  $1 \leq i \leq m$  and  $X$  is open and  $g = f$  and  $X = Y$ . Then the following statements are equivalent
- (i)  $f$  is partially differentiable on  $X$  w.r.t.  $i$  and  $f|_X^i$  is continuous on  $X$ ,

- (ii)  $g$  is partially differentiable on  $Y$  w.r.t.  $i$  and  $g|_Y^i$  is continuous on  $Y$ .
- (24) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $X$  be a subset of  $\mathcal{R}^m$ , and  $Y$  be a subset of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . Suppose  $X = Y$  and  $X$  is open and  $f = g$ . Then for every  $i$  such that  $1 \leq i \leq m$  holds  $f$  is partially differentiable on  $X$  w.r.t.  $i$  and  $f|_X^i$  is continuous on  $X$  if and only if  $g$  is differentiable on  $Y$  and  $g'|_Y$  is continuous on  $Y$ .
- (25) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $X$  be a subset of  $\mathcal{R}^m$ , and  $Y$  be a subset of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . Suppose  $X$  is open and  $X \subseteq \text{dom } f$  and  $g = f$  and  $X = Y$ . Then  $g$  is differentiable on  $Y$  and  $g'|_Y$  is continuous on  $Y$  if and only if the following conditions are satisfied:
  - (i)  $f$  is differentiable on  $X$ , and
  - (ii) for every element  $x_0$  of  $\mathcal{R}^m$  and for every real number  $r$  such that  $x_0 \in X$  and  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every element  $x_1$  of  $\mathcal{R}^m$  such that  $x_1 \in X$  and  $|x_1 - x_0| < s$  and for every element  $v$  of  $\mathcal{R}^m$  holds  $|f'(x_1)(v) - f'(x_0)(v)| \leq r \cdot |v|$ .
- (26) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . Suppose  $X$  is open and  $X \subseteq \text{dom } f$ . Then the following statements are equivalent
  - (i) for every element  $i$  of  $\mathbb{N}$  such that  $1 \leq i \leq m$  holds  $f$  is partially differentiable on  $X$  w.r.t.  $i$  and  $f|_X^i$  is continuous on  $X$ ,
  - (ii)  $f$  is differentiable on  $X$  and for every element  $x_0$  of  $\mathcal{R}^m$  and for every real number  $r$  such that  $x_0 \in X$  and  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every element  $x_1$  of  $\mathcal{R}^m$  such that  $x_1 \in X$  and  $|x_1 - x_0| < s$  and for every element  $v$  of  $\mathcal{R}^m$  holds  $|f'(x_1)(v) - f'(x_0)(v)| \leq r \cdot |v|$ .
- (27) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$  and  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ . If  $f = g$  and  $f$  is differentiable on  $Z$ , then  $f'|_Z = g'|_Z$ .
- (28) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\langle \mathcal{E}^n, \|\cdot\| \rangle$ ,  $X$  be a subset of  $\mathcal{R}^m$ , and  $Y$  be a subset of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$ . Suppose  $X = Y$  and  $X$  is open and  $f = g$ . Then for every element  $i$  of  $\mathbb{N}$  such that  $1 \leq i \leq m$  holds  $f$  is partially differentiable on  $X$  w.r.t.  $i$  and  $f|_X^i$  is continuous on  $X$  if and only if  $f$  is differentiable on  $X$  and  $g'|_Y$  is continuous on  $Y$ .
- (29) Let  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathcal{R}^n$  and  $x$  be an element of  $\mathcal{R}^m$ . Suppose  $f$  is continuous in  $x$  and  $g$  is continuous in  $x$ . Then  $f + g$  is continuous in  $x$  and  $f - g$  is continuous in  $x$ .
- (30) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ ,  $x$  be an element of  $\mathcal{R}^m$ , and  $r$  be a real number. If  $f$  is continuous in  $x$ , then  $r \cdot f$  is continuous in  $x$ .

- (31) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$  and  $x$  be an element of  $\mathcal{R}^m$ . If  $f$  is continuous in  $x$ , then  $-f$  is continuous in  $x$ .
- (32) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^n$  and  $x$  be an element of  $\mathcal{R}^m$ . If  $f$  is continuous in  $x$ , then  $|f|$  is continuous in  $x$ .
- (33) Let  $Z$  be a set and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . Suppose  $f$  is continuous on  $Z$  and  $g$  is continuous on  $Z$ . Then  $f + g$  is continuous on  $Z$  and  $f - g$  is continuous on  $Z$ .
- (34) Let  $r$  be a real number and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathcal{R}^n$ . If  $f$  is continuous on  $Z$ , then  $r \cdot f$  is continuous on  $Z$ .
- (35) For all partial functions  $f, g$  from  $\mathcal{R}^m$  to  $\mathcal{R}^n$  such that  $f$  is continuous on  $Z$  holds  $-f$  is continuous on  $Z$ .
- (36) Let  $f$  be a partial function from  $\mathcal{R}^i$  to  $\mathbb{R}$  and  $x_0$  be an element of  $\mathcal{R}^i$ . Then  $f$  is continuous in  $x_0$  if and only if the following conditions are satisfied:
  - (i)  $x_0 \in \text{dom } f$ , and
  - (ii) for every real number  $r$  such that  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every element  $x$  of  $\mathcal{R}^i$  such that  $x \in \text{dom } f$  and  $|x - x_0| < s$  holds  $|f_x - f_{x_0}| < r$ .
- (37) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $x_0$  be an element of  $\mathcal{R}^m$ . Then  $f$  is continuous in  $x_0$  if and only if  $\langle f \rangle$  is continuous in  $x_0$ .
- (38) Let  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $x_0$  be an element of  $\mathcal{R}^m$ . Suppose  $f$  is continuous in  $x_0$  and  $g$  is continuous in  $x_0$ . Then  $f + g$  is continuous in  $x_0$  and  $f - g$  is continuous in  $x_0$ .
- (39) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ ,  $x_0$  be an element of  $\mathcal{R}^m$ , and  $r$  be a real number. If  $f$  is continuous in  $x_0$ , then  $r \cdot f$  is continuous in  $x_0$ .
- (40) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $x_0$  be an element of  $\mathcal{R}^m$ . If  $f$  is continuous in  $x_0$ , then  $|f|$  is continuous in  $x_0$ .
- (41) Let  $f, g$  be partial functions from  $\mathcal{R}^i$  to  $\mathbb{R}$  and  $x$  be an element of  $\mathcal{R}^i$ . If  $f$  is continuous in  $x$  and  $g$  is continuous in  $x$ , then  $f \cdot g$  is continuous in  $x$ .

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $Z$  be a set, and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . We say that  $f$  is continuous on  $Z$  if and only if:

- (Def. 2) For every element  $x_0$  of  $\mathcal{R}^m$  such that  $x_0 \in Z$  holds  $f|_Z$  is continuous in  $x_0$ .

We now state a number of propositions:

- (42) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\mathbb{R}$ . Suppose  $f = g$ . Then  $Z \subseteq \text{dom } f$  and  $f$  is continuous on  $Z$  if and only if  $g$  is continuous on  $Z$ .
- (43) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\mathbb{R}$ . Suppose  $f = g$  and  $Z \subseteq \text{dom } f$ . Then  $f$  is continuous on  $Z$

if and only if for every sequence  $s$  of  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  such that  $\text{rng } s \subseteq Z$  and  $s$  is convergent and  $\lim s \in Z$  holds  $g_*s$  is convergent and  $g_{\lim s} = \lim(g_*s)$ .

- (44) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $g$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^1$ . Suppose  $\langle f \rangle = g$ . Then  $Z \subseteq \text{dom } f$  and  $f$  is continuous on  $Z$  if and only if  $g$  is continuous on  $Z$ .
- (45) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $Z \subseteq \text{dom } f$ . Then  $f$  is continuous on  $Z$  if and only if for every element  $x_0$  of  $\mathcal{R}^m$  and for every real number  $r$  such that  $x_0 \in Z$  and  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every element  $x_1$  of  $\mathcal{R}^m$  such that  $x_1 \in Z$  and  $|x_1 - x_0| < s$  holds  $|f_{x_1} - f_{x_0}| < r$ .
- (46) Let  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $f$  is continuous on  $Z$  and  $g$  is continuous on  $Z$  and  $Z \subseteq \text{dom } f$  and  $Z \subseteq \text{dom } g$ . Then  $f + g$  is continuous on  $Z$  and  $f - g$  is continuous on  $Z$ .
- (47) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $r$  be a real number. If  $Z \subseteq \text{dom } f$  and  $f$  is continuous on  $Z$ , then  $r \cdot f$  is continuous on  $Z$ .
- (48) Let  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $f$  is continuous on  $Z$  and  $g$  is continuous on  $Z$  and  $Z \subseteq \text{dom } f$  and  $Z \subseteq \text{dom } g$ . Then  $f \cdot g$  is continuous on  $Z$ .
- (49) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $g$  be a partial function from  $\langle \mathcal{E}^m, \|\cdot\| \rangle$  to  $\mathbb{R}$ . Suppose  $f = g$ . Then  $Z \subseteq \text{dom } f$  and  $f$  is continuous on  $Z$  if and only if  $g$  is continuous on  $Z$ .
- (50) For all partial functions  $f, g$  from  $\mathcal{R}^m$  to  $\mathcal{R}^n$  such that  $f$  is continuous on  $Z$  holds  $|f|$  is continuous on  $Z$ .
- (51) Let  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $x$  be an element of  $\mathcal{R}^m$ . Suppose  $f$  is differentiable in  $x$  and  $g$  is differentiable in  $x$ . Then  $f + g$  is differentiable in  $x$  and  $(f + g)'(x) = f'(x) + g'(x)$  and  $f - g$  is differentiable in  $x$  and  $(f - g)'(x) = f'(x) - g'(x)$ .
- (52) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ ,  $r$  be a real number, and  $x$  be an element of  $\mathcal{R}^m$ . Suppose  $f$  is differentiable in  $x$ . Then  $r \cdot f$  is differentiable in  $x$  and  $(r \cdot f)'(x) = r \cdot f'(x)$ .

Let  $Z$  be a set, let  $m$  be a non empty element of  $\mathbb{N}$ , and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . We say that  $f$  is differentiable on  $Z$  if and only if:

- (Def. 3) For every element  $x$  of  $\mathcal{R}^m$  such that  $x \in Z$  holds  $f \upharpoonright Z$  is differentiable in  $x$ .

Next we state three propositions:

- (53) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $g$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^1$ . Suppose  $\langle f \rangle = g$ . Then  $Z \subseteq \text{dom } f$  and  $f$  is differentiable on  $Z$  if and only if  $g$  is differentiable on  $Z$ .
- (54) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $X \subseteq \text{dom } f$  and  $X$  is open. Then  $f$  is differentiable on  $X$  if and

only if for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $f$  is differentiable in  $x$ .

- (55) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . If  $X \subseteq \text{dom } f$  and  $f$  is differentiable on  $X$ , then  $X$  is open.

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $Z$  be a set, and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Let us assume that  $Z \subseteq \text{dom } f$ . The functor  $f'_{|Z}$  yields a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}^{\mathcal{R}^m}$  and is defined by:

- (Def. 4)  $\text{dom}(f'_{|Z}) = Z$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in Z$  holds  $(f'_{|Z})_x = f'(x)$ .

One can prove the following four propositions:

- (56) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ , and  $g$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}^1$ . Suppose  $\langle f \rangle = g$  and  $X \subseteq \text{dom } f$  and  $f$  is differentiable on  $X$ . Then  $g$  is differentiable on  $X$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $(f'_{|X})_x = \text{proj}(1, 1) \cdot (g'_{|X})_x$ .
- (57) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $X \subseteq \text{dom } f$  and  $X \subseteq \text{dom } g$  and  $f$  is differentiable on  $X$  and  $g$  is differentiable on  $X$ . Then  $f + g$  is differentiable on  $X$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((f + g)'_{|X})_x = (f'_{|X})_x + (g'_{|X})_x$ .
- (58) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $X \subseteq \text{dom } f$  and  $X \subseteq \text{dom } g$  and  $f$  is differentiable on  $X$  and  $g$  is differentiable on  $X$ . Then  $f - g$  is differentiable on  $X$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((f - g)'_{|X})_x = (f'_{|X})_x - (g'_{|X})_x$ .
- (59) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ , and  $r$  be a real number. Suppose  $X \subseteq \text{dom } f$  and  $f$  is differentiable on  $X$ . Then  $r \cdot f$  is differentiable on  $X$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((r \cdot f)'_{|X})_x = r \cdot (f'_{|X})_x$ .

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $Z$  be a set, let  $i$  be an element of  $\mathbb{N}$ , and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . We say that  $f$  is partially differentiable on  $Z$  w.r.t.  $i$  if and only if:

- (Def. 5)  $Z \subseteq \text{dom } f$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in Z$  holds  $f|Z$  is partially differentiable in  $x$  w.r.t.  $i$ .

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $Z$  be a set, let  $i$  be an element of  $\mathbb{N}$ , and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Let us assume that  $f$  is partially differentiable on  $Z$  w.r.t.  $i$ . The functor  $f|^i Z$  yields a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and is defined as follows:

- (Def. 6)  $\text{dom}(f|^i Z) = Z$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in Z$  holds  $(f|^i Z)_x = \text{partdiff}(f, x, i)$ .

Next we state several propositions:

- (60) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $X$  is open and  $1 \leq i \leq m$ . Then  $f$  is partially differentiable on  $X$

w.r.t.  $i$  if and only if  $X \subseteq \text{dom } f$  and for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $f$  is partially differentiable in  $x$  w.r.t.  $i$ .

- (61) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ , and  $g$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^1$ . Suppose  $\langle f \rangle = g$  and  $X$  is open and  $1 \leq i \leq m$ . Then  $f$  is partially differentiable on  $X$  w.r.t.  $i$  if and only if  $g$  is partially differentiable on  $X$  w.r.t.  $i$ .
- (62) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ , and  $g$  be a partial function from  $\mathcal{R}^m$  to  $\mathcal{R}^1$ . Suppose  $\langle f \rangle = g$  and  $X$  is open and  $1 \leq i \leq m$  and  $f$  is partially differentiable on  $X$  w.r.t.  $i$ . Then  $f|X$  is continuous on  $X$  if and only if  $g|X$  is continuous on  $X$ .
- (63) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $X$  is open and  $X \subseteq \text{dom } f$ . Then the following statements are equivalent
- (i) for every element  $i$  of  $\mathbb{N}$  such that  $1 \leq i \leq m$  holds  $f$  is partially differentiable on  $X$  w.r.t.  $i$  and  $f|X$  is continuous on  $X$ ,
  - (ii)  $f$  is differentiable on  $X$  and for every element  $x_0$  of  $\mathcal{R}^m$  and for every real number  $r$  such that  $x_0 \in X$  and  $0 < r$  there exists a real number  $s$  such that  $0 < s$  and for every element  $x_1$  of  $\mathcal{R}^m$  such that  $x_1 \in X$  and  $|x_1 - x_0| < s$  and for every element  $v$  of  $\mathcal{R}^m$  holds  $|f'(x_1)(v) - f'(x_0)(v)| \leq r \cdot |v|$ .
- (64) Let  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $x$  be an element of  $\mathcal{R}^m$ . Suppose  $f$  is partially differentiable in  $x$  w.r.t.  $i$  and  $g$  is partially differentiable in  $x$  w.r.t.  $i$ . Then  $f \cdot g$  is partially differentiable in  $x$  w.r.t.  $i$  and  $\text{partdiff}(f \cdot g, x, i) = \text{partdiff}(f, x, i) \cdot g(x) + f(x) \cdot \text{partdiff}(g, x, i)$ .
- (65) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that
- (i)  $X$  is open,
  - (ii)  $1 \leq i$ ,
  - (iii)  $i \leq m$ ,
  - (iv)  $f$  is partially differentiable on  $X$  w.r.t.  $i$ , and
  - (v)  $g$  is partially differentiable on  $X$  w.r.t.  $i$ .
- Then
- (vi)  $f + g$  is partially differentiable on  $X$  w.r.t.  $i$ ,
  - (vii)  $(f + g)|X = (f|X) + (g|X)$ , and
  - (viii) for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((f + g)|X)_x = \text{partdiff}(f, x, i) + \text{partdiff}(g, x, i)$ .
- (66) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that
- (i)  $X$  is open,
  - (ii)  $1 \leq i$ ,
  - (iii)  $i \leq m$ ,

- (iv)  $f$  is partially differentiable on  $X$  w.r.t.  $i$ , and
- (v)  $g$  is partially differentiable on  $X$  w.r.t.  $i$ .

Then

- (vi)  $f - g$  is partially differentiable on  $X$  w.r.t.  $i$ ,
  - (vii)  $(f - g) \upharpoonright^i X = (f \upharpoonright^i X) - (g \upharpoonright^i X)$ , and
  - (viii) for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $((f - g) \upharpoonright^i X)_x = \text{partdiff}(f, x, i) - \text{partdiff}(g, x, i)$ .
- (67) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $r$  be a real number, and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $X$  is open and  $1 \leq i \leq m$  and  $f$  is partially differentiable on  $X$  w.r.t.  $i$ . Then
- (i)  $r \cdot f$  is partially differentiable on  $X$  w.r.t.  $i$ ,
  - (ii)  $r \cdot f \upharpoonright^i X = r \cdot (f \upharpoonright^i X)$ , and
  - (iii) for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $(r \cdot f \upharpoonright^i X)_x = r \cdot \text{partdiff}(f, x, i)$ .

- (68) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that

- (i)  $X$  is open,
- (ii)  $1 \leq i$ ,
- (iii)  $i \leq m$ ,
- (iv)  $f$  is partially differentiable on  $X$  w.r.t.  $i$ , and
- (v)  $g$  is partially differentiable on  $X$  w.r.t.  $i$ .

Then

- (vi)  $f \cdot g$  is partially differentiable on  $X$  w.r.t.  $i$ ,
- (vii)  $f \cdot g \upharpoonright^i X = (f \upharpoonright^i X) \cdot g + f \cdot (g \upharpoonright^i X)$ , and
- (viii) for every element  $x$  of  $\mathcal{R}^m$  such that  $x \in X$  holds  $(f \cdot g \upharpoonright^i X)_x = \text{partdiff}(f, x, i) \cdot g(x) + f(x) \cdot \text{partdiff}(g, x, i)$ .

### 3. HIGHER-ORDER PARTIAL DIFFERENTIATION

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $Z$  be a set, let  $I$  be a finite sequence of elements of  $\mathbb{N}$ , and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . The functor  $\text{PartDiffSeq}(f, Z, I)$  yielding a sequence of partial functions from  $\mathcal{R}^m$  into  $\mathbb{R}$  is defined by:

- (Def. 7)  $(\text{PartDiffSeq}(f, Z, I))(0) = f$  and for every natural number  $i$  holds  $(\text{PartDiffSeq}(f, Z, I))(i+1) = (\text{PartDiffSeq}(f, Z, I))(i) \upharpoonright^{I_{i+1}} Z$ .

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $Z$  be a set, let  $I$  be a finite sequence of elements of  $\mathbb{N}$ , and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . We say that  $f$  is partially differentiable on  $Z$  w.r.t.  $I$  if and only if:

- (Def. 8) For every element  $i$  of  $\mathbb{N}$  such that  $i \leq \text{len } I - 1$  holds  $(\text{PartDiffSeq}(f, Z, I))(i)$  is partially differentiable on  $Z$  w.r.t.  $I_{i+1}$ .

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $Z$  be a set, let  $I$  be a finite sequence of elements of  $\mathbb{N}$ , and let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . The functor  $f|{}^I Z$  yielding a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  is defined by:

(Def. 9)  $f|{}^I Z = (\text{PartDiffSeq}(f, Z, I))(\text{len } I)$ .

The following propositions are true:

(69) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ , and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that

- (i)  $X$  is open,
- (ii)  $\text{rng } I \subseteq \text{Seg } m$ ,
- (iii)  $f$  is partially differentiable on  $X$  w.r.t.  $I$ , and
- (iv)  $g$  is partially differentiable on  $X$  w.r.t.  $I$ .

Let given  $i$ . Suppose  $i \leq \text{len } I - 1$ . Then  $(\text{PartDiffSeq}(f + g, X, I))(i)$  is partially differentiable on  $X$  w.r.t.  $I_{i+1}$  and  $(\text{PartDiffSeq}(f + g, X, I))(i) = (\text{PartDiffSeq}(f, X, I))(i) + (\text{PartDiffSeq}(g, X, I))(i)$ .

(70) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ , and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that

- (i)  $X$  is open,
- (ii)  $\text{rng } I \subseteq \text{Seg } m$ ,
- (iii)  $f$  is partially differentiable on  $X$  w.r.t.  $I$ , and
- (iv)  $g$  is partially differentiable on  $X$  w.r.t.  $I$ .

Then  $f + g$  is partially differentiable on  $X$  w.r.t.  $I$  and  $(f + g)|{}^I X = (f|{}^I X) + (g|{}^I X)$ .

(71) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ , and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that

- (i)  $X$  is open,
- (ii)  $\text{rng } I \subseteq \text{Seg } m$ ,
- (iii)  $f$  is partially differentiable on  $X$  w.r.t.  $I$ , and
- (iv)  $g$  is partially differentiable on  $X$  w.r.t.  $I$ .

Let given  $i$ . Suppose  $i \leq \text{len } I - 1$ . Then  $(\text{PartDiffSeq}(f - g, X, I))(i)$  is partially differentiable on  $X$  w.r.t.  $I_{i+1}$  and  $(\text{PartDiffSeq}(f - g, X, I))(i) = (\text{PartDiffSeq}(f, X, I))(i) - (\text{PartDiffSeq}(g, X, I))(i)$ .

(72) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ , and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that

- (i)  $X$  is open,
- (ii)  $\text{rng } I \subseteq \text{Seg } m$ ,
- (iii)  $f$  is partially differentiable on  $X$  w.r.t.  $I$ , and
- (iv)  $g$  is partially differentiable on  $X$  w.r.t.  $I$ .

Then  $f - g$  is partially differentiable on  $X$  w.r.t.  $I$  and  $(f - g)|{}^I X = (f|{}^I X) - (g|{}^I X)$ .

(73) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $r$  be a real number,  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ , and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ .

Suppose  $X$  is open and  $\text{rng } I \subseteq \text{Seg } m$  and  $f$  is partially differentiable on  $X$  w.r.t.  $I$ . Let given  $i$ . Suppose  $i \leq \text{len } I - 1$ . Then  $(\text{PartDiffSeq}(r \cdot f, X, I))(i)$  is partially differentiable on  $X$  w.r.t.  $I_{i+1}$  and  $(\text{PartDiffSeq}(r \cdot f, X, I))(i) = r \cdot (\text{PartDiffSeq}(f, X, I))(i)$ .

- (74) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $r$  be a real number,  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ , and  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $X$  is open and  $\text{rng } I \subseteq \text{Seg } m$  and  $f$  is partially differentiable on  $X$  w.r.t.  $I$ . Then  $r \cdot f$  is partially differentiable on  $X$  w.r.t.  $I$  and  $r \cdot f \upharpoonright^I X = r \cdot (f \upharpoonright^I X)$ .

Let  $m$  be a non empty element of  $\mathbb{N}$ , let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ , let  $k$  be an element of  $\mathbb{N}$ , and let  $Z$  be a set. We say that  $f$  is partial differentiable up to order  $k$  and  $Z$  if and only if the condition (Def. 10) is satisfied.

- (Def. 10) Let  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ . If  $\text{len } I \leq k$  and  $\text{rng } I \subseteq \text{Seg } m$ , then  $f$  is partially differentiable on  $Z$  w.r.t.  $I$ .

The following proposition is true

- (75) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $I, G$  be non empty finite sequences of elements of  $\mathbb{N}$ . Then  $f$  is partially differentiable on  $Z$  w.r.t.  $G \cap I$  if and only if  $f$  is partially differentiable on  $Z$  w.r.t.  $G$  and  $f \upharpoonright^G Z$  is partially differentiable on  $Z$  w.r.t.  $I$ .

One can prove the following propositions:

- (76) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Then  $f$  is partially differentiable on  $Z$  w.r.t.  $\langle i \rangle$  if and only if  $f$  is partially differentiable on  $Z$  w.r.t.  $i$ .
- (77) For every partial function  $f$  from  $\mathcal{R}^m$  to  $\mathbb{R}$  holds  $f \upharpoonright^{\langle i \rangle} Z = f \upharpoonright^i Z$ .
- (78) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$  and  $I$  be a non empty finite sequence of elements of  $\mathbb{N}$ . Suppose  $f$  is partial differentiable up to order  $i + j$  and  $Z$  and  $\text{rng } I \subseteq \text{Seg } m$  and  $\text{len } I = j$ . Then  $f \upharpoonright^I Z$  is partial differentiable up to order  $i$  and  $Z$ .
- (79) Let  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $f$  is partial differentiable up to order  $i$  and  $Z$  and  $j \leq i$ . Then  $f$  is partial differentiable up to order  $j$  and  $Z$ .
- (80) Let  $X$  be a subset of  $\mathcal{R}^m$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose that
- (i)  $X$  is open,
  - (ii)  $f$  is partial differentiable up to order  $i$  and  $X$ , and
  - (iii)  $g$  is partial differentiable up to order  $i$  and  $X$ .
- Then  $f + g$  is partial differentiable up to order  $i$  and  $X$  and  $f - g$  is partial differentiable up to order  $i$  and  $X$ .
- (81) Let  $X$  be a subset of  $\mathcal{R}^m$ ,  $f$  be a partial function from  $\mathcal{R}^m$  to  $\mathbb{R}$ , and  $r$  be a real number. Suppose  $X$  is open and  $f$  is partial differentiable up to

order  $i$  and  $X$ . Then  $r \cdot f$  is partial differentiable up to order  $i$  and  $X$ .

- (82) Let  $X$  be a subset of  $\mathcal{R}^m$ . Suppose  $X$  is open. Let  $i$  be an element of  $\mathbb{N}$  and  $f, g$  be partial functions from  $\mathcal{R}^m$  to  $\mathbb{R}$ . Suppose  $f$  is partial differentiable up to order  $i$  and  $X$  and  $g$  is partial differentiable up to order  $i$  and  $X$ . Then  $f \cdot g$  is partial differentiable up to order  $i$  and  $X$ .

## REFERENCES

- [1] Grzegorz Bancerek. The ordinal numbers. *Formalized Mathematics*, 1(1):91–96, 1990.
- [2] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [3] Czesław Byliński. The complex numbers. *Formalized Mathematics*, 1(3):507–513, 1990.
- [4] Czesław Byliński. Finite sequences and tuples of elements of a non-empty sets. *Formalized Mathematics*, 1(3):529–536, 1990.
- [5] Czesław Byliński. Functions from a set to a set. *Formalized Mathematics*, 1(1):153–164, 1990.
- [6] Czesław Byliński. Partial functions. *Formalized Mathematics*, 1(2):357–367, 1990.
- [7] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [8] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [9] Noboru Endou and Yasunari Shidama. Completeness of the real Euclidean space. *Formalized Mathematics*, 13(4):577–580, 2005.
- [10] Noboru Endou, Yasunari Shidama, and Keiichi Miyajima. Partial differentiation on normed linear spaces  $\mathcal{R}^n$ . *Formalized Mathematics*, 15(2):65–72, 2007, doi:10.2478/v10037-007-0008-5.
- [11] Krzysztof Hryniewiecki. Basic properties of real numbers. *Formalized Mathematics*, 1(1):35–40, 1990.
- [12] Takao Inoué, Noboru Endou, and Yasunari Shidama. Differentiation of vector-valued functions on  $n$ -dimensional real normed linear spaces. *Formalized Mathematics*, 18(4):207–212, 2010, doi: 10.2478/v10037-010-0025-7.
- [13] Takao Inoué, Adam Naumowicz, Noboru Endou, and Yasunari Shidama. Partial differentiation of vector-valued functions on  $n$ -dimensional real normed linear spaces. *Formalized Mathematics*, 19(1):1–9, 2011, doi: 10.2478/v10037-011-0001-x.
- [14] Jarosław Kotowicz. Real sequences and basic operations on them. *Formalized Mathematics*, 1(2):269–272, 1990.
- [15] Keiichi Miyajima and Yasunari Shidama. Riemann integral of functions from  $\mathbb{R}$  into  $\mathcal{R}^n$ . *Formalized Mathematics*, 17(2):179–185, 2009, doi: 10.2478/v10037-009-0021-y.
- [16] Keiko Narita, Artur Kornilowicz, and Yasunari Shidama. More on the continuity of real functions. *Formalized Mathematics*, 19(4):233–239, 2011, doi: 10.2478/v10037-011-0032-3.
- [17] Takaya Nishiyama, Keiji Ohkubo, and Yasunari Shidama. The continuous functions on normed linear spaces. *Formalized Mathematics*, 12(3):269–275, 2004.
- [18] Beata Padlewska and Agata Darmochwał. Topological spaces and continuous functions. *Formalized Mathematics*, 1(1):223–230, 1990.
- [19] Beata Perkowska. Functional sequence from a domain to a domain. *Formalized Mathematics*, 3(1):17–21, 1992.
- [20] Jan Popiołek. Real normed space. *Formalized Mathematics*, 2(1):111–115, 1991.
- [21] Yasunari Shidama. Banach space of bounded linear operators. *Formalized Mathematics*, 12(1):39–48, 2004.
- [22] Zinaida Trybulec. Properties of subsets. *Formalized Mathematics*, 1(1):67–71, 1990.
- [23] Edmund Woronowicz. Relations defined on sets. *Formalized Mathematics*, 1(1):181–186, 1990.

Received November 20, 2011