

# Group of Homography in Real Projective Plane

Roland Coghetto  
Rue de la Brasserie 5  
7100 La Louvière, Belgium

**Summary.** Using the Mizar system [2], we formalized that homographies of the projective real plane (as defined in [5]), form a group.

Then, we prove that, using the notations of Borsuk and Szmielew in [3]

“Consider in space  $\mathbb{RP}^2$  points  $P_1, P_2, P_3, P_4$  of which three points are not collinear and points  $Q_1, Q_2, Q_3, Q_4$  each three points of which are also not collinear. There exists one homography  $h$  of space  $\mathbb{RP}^2$  such that  $h(P_i) = Q_i$  for  $i = 1, 2, 3, 4$ .”

(Existence Statement 52 and Existence Statement 53) [3]. Or, using notations of Richter [11]

“Let  $[a], [b], [c], [d]$  in  $\mathbb{RP}^2$  be four points of which no three are collinear and let  $[a'], [b'], [c'], [d']$  in  $\mathbb{RP}^2$  be another four points of which no three are collinear, then there exists a  $3 \times 3$  matrix  $M$  such that  $[Ma] = [a'], [Mb] = [b'], [Mc] = [c'],$  and  $[Md] = [d']$ ”

Makarios has formalized the same results in Isabelle/Isar (the collineations form a group, lemma statement52-existence and lemma statement 53-existence) and published it in Archive of Formal Proofs<sup>1</sup> [10], [9].

MSC: 51N15 03B35

Keywords: projectivity; projective transformation; real projective plane; group of homography

MML identifier: ANPROJ\_9, version: 8.1.05 5.40.1289

---

<sup>1</sup>[http://isa-afp.org/entries/Tarskis\\_Geometry.shtml](http://isa-afp.org/entries/Tarskis_Geometry.shtml)

## 1. PRELIMINARIES

From now on  $i, n$  denote natural numbers,  $r$  denotes a real number,  $r_1$  denotes an element of  $\mathbb{R}_F$ ,  $a, b, c$  denote non zero elements of  $\mathbb{R}_F$ ,  $u, v$  denote elements of  $\mathcal{E}_T^3$ ,  $p_1$  denotes a finite sequence of elements of  $\mathbb{R}^1$ ,  $p_3, u_4$  denote finite sequences of elements of  $\mathbb{R}_F$ ,  $N$  denotes a square matrix over  $\mathbb{R}_F$  of dimension 3,  $K$  denotes a field, and  $k$  denotes an element of  $K$ .

Now we state the propositions:

- (1)  $I_{\mathbb{R}_F}^{3 \times 3} = \langle \langle 1, 0, 0 \rangle, \langle 0, 1, 0 \rangle, \langle 0, 0, 1 \rangle \rangle$ .
- (2)  $r_1 \cdot N = r_1 \cdot I_{\mathbb{R}_F}^{3 \times 3} \cdot N$ .
- (3) If  $r \neq 0$  and  $u$  is not zero, then  $r \cdot u$  is not zero.

PROOF:  $r \cdot u \neq 0_{\mathcal{E}_T^3}$  by [4, (52), (49)].  $\square$

Let us consider elements  $a_{11}, a_{12}, a_{13}, a_{21}, a_{22}, a_{23}, a_{31}, a_{32}, a_{33}$  of  $\mathbb{R}_F$  and a square matrix  $A$  over  $\mathbb{R}_F$  of dimension 3. Now we state the propositions:

- (4) Suppose  $A = \langle \langle a_{11}, a_{12}, a_{13} \rangle, \langle a_{21}, a_{22}, a_{23} \rangle, \langle a_{31}, a_{32}, a_{33} \rangle \rangle$ . Then
  - (i)  $\text{Line}(A, 1) = \langle a_{11}, a_{12}, a_{13} \rangle$ , and
  - (ii)  $\text{Line}(A, 2) = \langle a_{21}, a_{22}, a_{23} \rangle$ , and
  - (iii)  $\text{Line}(A, 3) = \langle a_{31}, a_{32}, a_{33} \rangle$ .
- (5) Suppose  $A = \langle \langle a_{11}, a_{12}, a_{13} \rangle, \langle a_{21}, a_{22}, a_{23} \rangle, \langle a_{31}, a_{32}, a_{33} \rangle \rangle$ . Then
  - (i)  $A_{\square, 1} = \langle a_{11}, a_{21}, a_{31} \rangle$ , and
  - (ii)  $A_{\square, 2} = \langle a_{12}, a_{22}, a_{32} \rangle$ , and
  - (iii)  $A_{\square, 3} = \langle a_{13}, a_{23}, a_{33} \rangle$ .

The theorem is a consequence of (4).

- (6) Let us consider elements  $a_{11}, a_{12}, a_{13}, a_{21}, a_{22}, a_{23}, a_{31}, a_{32}, a_{33}, b_{11}, b_{12}, b_{13}, b_{21}, b_{22}, b_{23}, b_{31}, b_{32}, b_{33}$  of  $\mathbb{R}_F$ , and square matrices  $A, B$  over  $\mathbb{R}_F$  of dimension 3. Suppose  $A = \langle \langle a_{11}, a_{12}, a_{13} \rangle, \langle a_{21}, a_{22}, a_{23} \rangle, \langle a_{31}, a_{32}, a_{33} \rangle \rangle$  and  $B = \langle \langle b_{11}, b_{12}, b_{13} \rangle, \langle b_{21}, b_{22}, b_{23} \rangle, \langle b_{31}, b_{32}, b_{33} \rangle \rangle$ . Then  $A \cdot B = \langle \langle a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}, a_{11} \cdot b_{12} + a_{12} \cdot b_{22} + a_{13} \cdot b_{32}, a_{11} \cdot b_{13} + a_{12} \cdot b_{23} + a_{13} \cdot b_{33} \rangle, \langle a_{21} \cdot b_{11} + a_{22} \cdot b_{21} + a_{23} \cdot b_{31}, a_{21} \cdot b_{12} + a_{22} \cdot b_{22} + a_{23} \cdot b_{32}, a_{21} \cdot b_{13} + a_{22} \cdot b_{23} + a_{23} \cdot b_{33} \rangle, \langle a_{31} \cdot b_{11} + a_{32} \cdot b_{21} + a_{33} \cdot b_{31}, a_{31} \cdot b_{12} + a_{32} \cdot b_{22} + a_{33} \cdot b_{32}, a_{31} \cdot b_{13} + a_{32} \cdot b_{23} + a_{33} \cdot b_{33} \rangle \rangle$ . The theorem is a consequence of (4) and (5).
- (7) Let us consider elements  $a_{11}, a_{12}, a_{13}, a_{21}, a_{22}, a_{23}, a_{31}, a_{32}, a_{33}, b_1, b_2, b_3$  of  $\mathbb{R}_F$ , a matrix  $A$  over  $\mathbb{R}_F$  of dimension  $3 \times 3$ , and a matrix  $B$  over  $\mathbb{R}_F$  of dimension  $3 \times 1$ . Suppose  $A = \langle \langle a_{11}, a_{12}, a_{13} \rangle, \langle a_{21}, a_{22}, a_{23} \rangle, \langle a_{31}, a_{32}, a_{33} \rangle \rangle$  and  $B = \langle \langle b_1 \rangle, \langle b_2 \rangle, \langle b_3 \rangle \rangle$ . Then  $A \cdot B = \langle \langle a_{11} \cdot b_1 + a_{12} \cdot b_2 + a_{13} \cdot b_3 \rangle, \langle a_{21} \cdot b_1 + a_{22} \cdot b_2 + a_{23} \cdot b_3 \rangle, \langle a_{31} \cdot b_1 + a_{32} \cdot b_2 + a_{33} \cdot b_3 \rangle \rangle$ .

$\langle a_{21} \cdot b_1 + a_{22} \cdot b_2 + a_{23} \cdot b_3 \rangle, \langle a_{31} \cdot b_1 + a_{32} \cdot b_2 + a_{33} \cdot b_3 \rangle$ . The theorem is a consequence of (4).

- (8) Let us consider non zero elements  $a, b, c$  of  $\mathbb{R}_F$ , and square matrices  $M_1, M_2$  over  $\mathbb{R}_F$  of dimension 3. Suppose  $M_1 = \langle \langle a, 0, 0 \rangle, \langle 0, b, 0 \rangle, \langle 0, 0, c \rangle \rangle$  and  $M_2 = \langle \langle \frac{1}{a}, 0, 0 \rangle, \langle 0, \frac{1}{b}, 0 \rangle, \langle 0, 0, \frac{1}{c} \rangle \rangle$ . Then

- (i)  $M_1 \cdot M_2 = I_{\mathbb{R}_F}^{3 \times 3}$ , and  
(ii)  $M_2 \cdot M_1 = I_{\mathbb{R}_F}^{3 \times 3}$ .

The theorem is a consequence of (1).

- (9) Let us consider non zero elements  $a, b, c$  of  $\mathbb{R}_F$ . Then  $\langle \langle a, 0, 0 \rangle, \langle 0, b, 0 \rangle, \langle 0, 0, c \rangle \rangle$  is an invertible square matrix over  $\mathbb{R}_F$  of dimension 3. The theorem is a consequence of (8).

- (10) (i)  $[1, 0, 0]$  is not zero, and

(ii)  $[0, 1, 0]$  is not zero, and

(iii)  $[0, 0, 1]$  is not zero, and

(iv)  $[1, 1, 1]$  is not zero.

- (11) (i)  $[1, 0, 0] \neq 0_{\mathcal{E}_T^3}$ , and

(ii)  $[0, 1, 0] \neq 0_{\mathcal{E}_T^3}$ , and

(iii)  $[0, 0, 1] \neq 0_{\mathcal{E}_T^3}$ , and

(iv)  $[1, 1, 1] \neq 0_{\mathcal{E}_T^3}$ .

PROOF:  $[1, 0, 0] \neq [0, 0, 0]$  by [7, (2)].  $[0, 1, 0] \neq [0, 0, 0]$  by [7, (2)].  $[0, 0, 1] \neq [0, 0, 0]$  by [7, (2)].  $[1, 1, 1] \neq [0, 0, 0]$  by [7, (2)].  $\square$

- (12) (i)  $e_1 \neq 0_{\mathcal{E}_T^3}$ , and

(ii)  $e_2 \neq 0_{\mathcal{E}_T^3}$ , and

(iii)  $e_3 \neq 0_{\mathcal{E}_T^3}$ .

PROOF:  $[1, 0, 0] \neq [0, 0, 0]$  by [7, (2)].  $[0, 1, 0] \neq [0, 0, 0]$  by [7, (2)].  $[0, 0, 1] \neq [0, 0, 0]$  by [7, (2)].  $\square$

Let  $n$  be a natural number. Note that  $I_{\mathbb{R}_F}^{n \times n}$  is invertible.

Let  $M$  be an invertible square matrix over  $\mathbb{R}_F$  of dimension  $n$ . One can verify that  $M^\sim$  is invertible.

Let  $K$  be a field and  $N_1, N_2$  be invertible square matrices over  $K$  of dimension  $n$ . One can check that  $N_1 \cdot N_2$  is invertible.

## 2. GROUP OF HOMOGRAPHY

From now on  $N, N_1, N_2$  denote invertible square matrices over  $\mathbb{R}_F$  of dimension 3 and  $P, P_1, P_2, P_3$  denote points of the projective space over  $\mathcal{E}_T^3$ .

Now we state the propositions:

- (13) (The homography of  $N_1$ )(the homography of  $N_2$ )( $P$ ) =  
(the homography of  $N_1 \cdot N_2$ )( $P$ ).

PROOF: Consider  $u_{12}, v_{12}$  being elements of  $\mathcal{E}_T^3$ ,  $u_8$  being a finite sequence of elements of  $\mathbb{R}_F$ ,  $p_{12}$  being a finite sequence of elements of  $\mathbb{R}^1$  such that  $P$  = the direction of  $u_{12}$  and  $u_{12}$  is not zero and  $u_{12} = u_8$  and  $p_{12} = N_1 \cdot N_2 \cdot u_8$  and  $v_{12} = \text{M2F}(p_{12})$  and  $v_{12}$  is not zero and (the homography of  $N_1 \cdot N_2$ )( $P$ ) = the direction of  $v_{12}$ . Consider  $u_2, v_2$  being elements of  $\mathcal{E}_T^3$ ,  $u_6$  being a finite sequence of elements of  $\mathbb{R}_F$ ,  $p_2$  being a finite sequence of elements of  $\mathbb{R}^1$  such that  $P$  = the direction of  $u_2$  and  $u_2$  is not zero and  $u_2 = u_6$  and  $p_2 = N_2 \cdot u_6$  and  $v_2 = \text{M2F}(p_2)$  and  $v_2$  is not zero and (the homography of  $N_2$ )( $P$ ) = the direction of  $v_2$ . Consider  $u_1, v_1$  being elements of  $\mathcal{E}_T^3$ ,  $u_7$  being a finite sequence of elements of  $\mathbb{R}_F$ ,  $p_1$  being a finite sequence of elements of  $\mathbb{R}^1$  such that (the homography of  $N_2$ )( $P$ ) = the direction of  $u_1$  and  $u_1$  is not zero and  $u_1 = u_7$  and  $p_1 = N_1 \cdot u_7$  and  $v_1 = \text{M2F}(p_1)$  and  $v_1$  is not zero and (the homography of  $N_1$ )(the homography of  $N_2$ )( $P$ ) = the direction of  $v_1$ . Consider  $a$  being a real number such that  $a \neq 0$  and  $u_2 = a \cdot u_{12}$ . Consider  $b$  being a real number such that  $b \neq 0$  and  $u_1 = b \cdot v_2$ .  $v_1 = \langle (N_1 \cdot \langle u_7 \rangle^T)_{1,1}, (N_1 \cdot \langle u_7 \rangle^T)_{2,1}, (N_1 \cdot \langle u_7 \rangle^T)_{3,1} \rangle$  by [1, (1), (40)].  $v_2 = \langle (N_2 \cdot \langle u_6 \rangle^T)_{1,1}, (N_2 \cdot \langle u_6 \rangle^T)_{2,1}, (N_2 \cdot \langle u_6 \rangle^T)_{3,1} \rangle$  by [1, (1), (40)].  $v_{12} = \langle (N_1 \cdot N_2 \cdot \langle u_8 \rangle^T)_{1,1}, (N_1 \cdot N_2 \cdot \langle u_8 \rangle^T)_{2,1}, (N_1 \cdot N_2 \cdot \langle u_8 \rangle^T)_{3,1} \rangle$  by [1, (1), (40)]. Reconsider  $v_6 = v_2$  as a finite sequence of elements of  $\mathbb{R}_F$ . Reconsider  $i_4 = \frac{1}{b}$  as a real number.  $v_6 = i_4 \cdot u_1$  by [4, (49), (52)]. Reconsider  $l_{11} = \text{Line}(N_2, 1)(1)$ ,  $l_{12} = \text{Line}(N_2, 1)(2)$ ,  $l_{13} = \text{Line}(N_2, 1)(3)$ ,  $l_{21} = \text{Line}(N_2, 2)(1)$ ,  $l_{22} = \text{Line}(N_2, 2)(2)$ ,  $l_{23} = \text{Line}(N_2, 2)(3)$ ,  $l_{31} = \text{Line}(N_2, 3)(1)$ ,  $l_{32} = \text{Line}(N_2, 3)(2)$ ,  $l_{33} = \text{Line}(N_2, 3)(3)$  as an element of  $\mathbb{R}_F$ .  $N_{2\Box,1} = \langle l_{11}, l_{21}, l_{31} \rangle$  and  $N_{2\Box,2} = \langle l_{12}, l_{22}, l_{32} \rangle$  and  $N_{2\Box,3} = \langle l_{13}, l_{23}, l_{33} \rangle$  by [1, (1), (45)]. The direction of  $v_1$  = the direction of  $v_{12}$  by [5, (7)], [1, (45)], [5, (93)], [7, (8)].  $\square$

- (14) (The homography of  $I_{\mathbb{R}_F}^{3 \times 3}$ )( $P$ ) =  $P$ .

- (15) (i) (the homography of  $N$ )(the homography of  $N^\sim$ )( $P$ ) =  $P$ , and  
(ii) (the homography of  $N^\sim$ )(the homography of  $N$ )( $P$ ) =  $P$ .

The theorem is a consequence of (13) and (14).

- (16) If (the homography of  $N$ )( $P_1$ ) = (the homography of  $N$ )( $P_2$ ), then  $P_1 = P_2$ . The theorem is a consequence of (15).

- (17) Let us consider a non zero element  $a$  of  $\mathbb{R}_F$ . Suppose  $a \cdot I_{\mathbb{R}_F}^{3 \times 3} = N$ . Then (the homography of  $N$ )( $P$ ) =  $P$ .

The functor  $\text{EnsHomography3}$  yielding a set is defined by the term

- (Def. 1) the set of all the homography of  $N$  where  $N$  is an invertible square matrix over  $\mathbb{R}_F$  of dimension 3.

One can check that  $\text{EnsHomography3}$  is non empty.

Let  $h_1, h_2$  be elements of  $\text{EnsHomography3}$ . The functor  $h_1 \circ h_2$  yielding an element of  $\text{EnsHomography3}$  is defined by

- (Def. 2) there exist invertible square matrices  $N_1, N_2$  over  $\mathbb{R}_F$  of dimension 3 such that  $h_1 =$  the homography of  $N_1$  and  $h_2 =$  the homography of  $N_2$  and  $it =$  the homography of  $N_1 \cdot N_2$ .

Now we state the propositions:

- (18) Let us consider elements  $h_1, h_2$  of  $\text{EnsHomography3}$ . Suppose  $h_1 =$  the homography of  $N_1$  and  $h_2 =$  the homography of  $N_2$ . Then the homography of  $N_1 \cdot N_2 = h_1 \circ h_2$ . The theorem is a consequence of (13).
- (19) Let us consider elements  $x, y, z$  of  $\text{EnsHomography3}$ . Then  $(x \circ y) \circ z = x \circ (y \circ z)$ . The theorem is a consequence of (18).

The functor  $\text{BinOpHomography3}$  yielding a binary operation on  $\text{EnsHomography3}$  is defined by

- (Def. 3) for every elements  $h_1, h_2$  of  $\text{EnsHomography3}$ ,  $it(h_1, h_2) = h_1 \circ h_2$ .

The functor  $\text{GroupHomography3}$  yielding a strict multiplicative magma is defined by the term

- (Def. 4)  $\langle \text{EnsHomography3}, \text{BinOpHomography3} \rangle$ .

Note that  $\text{GroupHomography3}$  is non empty, associative, and group-like.

Now we state the propositions:

- (20)  $\mathbf{1}_{\text{GroupHomography3}} =$  the homography of  $I_{\mathbb{R}_F}^{3 \times 3}$ .
- (21) Let us consider elements  $h, g$  of  $\text{GroupHomography3}$ , and invertible square matrices  $N, N_{10}$  over  $\mathbb{R}_F$  of dimension 3. Suppose  $h =$  the homography of  $N$  and  $g =$  the homography of  $N_{10}$  and  $N_{10} = N^\smile$ . Then  $g = h^{-1}$ . The theorem is a consequence of (20).

### 3. MAIN RESULTS

The functors:  $\text{Dir100}$ ,  $\text{Dir010}$ ,  $\text{Dir001}$ , and  $\text{Dir111}$  yielding points of the projective space over  $\mathcal{E}_T^3$  are defined by terms

- (Def. 5) the direction of  $[1, 0, 0]$ ,
- (Def. 6) the direction of  $[0, 1, 0]$ ,

(Def. 7) the direction of  $[0, 0, 1]$ ,

(Def. 8) the direction of  $[1, 1, 1]$ ,

respectively. Now we state the proposition:

- (22) (i)  $\text{Dir100} \neq \text{Dir010}$ , and  
(ii)  $\text{Dir100} \neq \text{Dir001}$ , and  
(iii)  $\text{Dir100} \neq \text{Dir111}$ , and  
(iv)  $\text{Dir010} \neq \text{Dir001}$ , and  
(v)  $\text{Dir010} \neq \text{Dir111}$ , and  
(vi)  $\text{Dir001} \neq \text{Dir111}$ .

Let  $a$  be a non zero element of  $\mathbb{R}_F$ . Let us consider  $N$ . Note that  $a \cdot N$  is invertible as a square matrix over  $\mathbb{R}_F$  of dimension 3.

- (23) Let us consider a non zero element  $a$  of  $\mathbb{R}_F$ . Then (the homography of  $a \cdot N_1$ )( $P$ ) = (the homography of  $N_1$ )( $P$ ). The theorem is a consequence of (2), (13), and (17).
- (24) Suppose  $P_1, P_2$  and  $P_3$  are not collinear. Then there exists an invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3 such that

- (i) (the homography of  $N$ )( $P_1$ ) =  $\text{Dir100}$ , and  
(ii) (the homography of  $N$ )( $P_2$ ) =  $\text{Dir010}$ , and  
(iii) (the homography of  $N$ )( $P_3$ ) =  $\text{Dir001}$ .

PROOF: Consider  $u_1$  being an element of  $\mathcal{E}_T^3$  such that  $u_1$  is not zero and  $P_1$  = the direction of  $u_1$ . Consider  $u_2$  being an element of  $\mathcal{E}_T^3$  such that  $u_2$  is not zero and  $P_2$  = the direction of  $u_2$ . Consider  $u_3$  being an element of  $\mathcal{E}_T^3$  such that  $u_3$  is not zero and  $P_3$  = the direction of  $u_3$ . Reconsider  $p_3 = u_1, q_1 = u_2, r_2 = u_3$  as a finite sequence of elements of  $\mathbb{R}_F$ . Consider  $N$  being a square matrix over  $\mathbb{R}_F$  of dimension 3 such that  $N$  is invertible and  $N \cdot p_3 = \text{F2M}(e_1)$  and  $N \cdot q_1 = \text{F2M}(e_2)$  and  $N \cdot r_2 = \text{F2M}(e_3)$ . (The homography of  $N$ )( $P_1$ ) =  $\text{Dir100}$  by [8, (22), (1)], [6, (22)], [5, (75)]. (The homography of  $N$ )( $P_2$ ) =  $\text{Dir010}$  by [8, (22), (1)], [6, (22)], [5, (75)]. (The homography of  $N$ )( $P_3$ ) =  $\text{Dir001}$  by [8, (22), (1)], [6, (22)], [5, (75)].  $\square$

- (25) Let us consider non zero elements  $a, b, c$  of  $\mathbb{R}_F$ . Suppose  $N = \langle \langle a, 0, 0 \rangle, \langle 0, b, 0 \rangle, \langle 0, 0, c \rangle \rangle$ . Then
- (i) (the homography of  $N$ )( $\text{Dir100}$ ) =  $\text{Dir100}$ , and  
(ii) (the homography of  $N$ )( $\text{Dir010}$ ) =  $\text{Dir010}$ , and  
(iii) (the homography of  $N$ )( $\text{Dir001}$ ) =  $\text{Dir001}$ .

PROOF: (The homography of  $N$ )(Dir100) = Dir100 by (12), [8, (22), (1)], [7, (8), (2)]. (The homography of  $N$ )(Dir010) = Dir010 by (12), [8, (22), (1)], [7, (8), (2)]. (The homography of  $N$ )(Dir001) = Dir001 by (12), [8, (22), (1)], [7, (8), (2)].  $\square$

Let us consider a point  $P$  of the projective space over  $\mathcal{E}_T^3$ .

- (26) There exist elements  $a, b, c$  of  $\mathbb{R}_F$  such that
- (i)  $P$  = the direction of  $[a, b, c]$ , and
  - (ii)  $a \neq 0$  or  $b \neq 0$  or  $c \neq 0$ .
- (27) Suppose Dir100, Dir010 and  $P$  are not collinear and Dir100, Dir001 and  $P$  are not collinear and Dir010, Dir001 and  $P$  are not collinear. Then there exist non zero elements  $a, b, c$  of  $\mathbb{R}_F$  such that  $P$  = the direction of  $[a, b, c]$ . The theorem is a consequence of (26).
- (28) Let us consider non zero elements  $a, b, c, i_1, i_2, i_3$  of  $\mathbb{R}_F$ , a point  $P$  of the projective space over  $\mathcal{E}_T^3$ , and an invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3. Suppose  $P$  = the direction of  $[a, b, c]$  and  $i_1 = \frac{1}{a}$  and  $i_2 = \frac{1}{b}$  and  $i_3 = \frac{1}{c}$  and  $N = \langle \langle i_1, 0, 0 \rangle, \langle 0, i_2, 0 \rangle, \langle 0, 0, i_3 \rangle \rangle$ . Then (the homography of  $N$ )( $P$ ) = the direction of  $[1, 1, 1]$ .
- (29) Let us consider a point  $P$  of the projective space over  $\mathcal{E}_T^3$ . Suppose Dir100, Dir010 and  $P$  are not collinear and Dir100, Dir001 and  $P$  are not collinear and Dir010, Dir001 and  $P$  are not collinear. Then there exist non zero elements  $a, b, c$  of  $\mathbb{R}_F$  such that for every invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3 such that  $N = \langle \langle a, 0, 0 \rangle, \langle 0, b, 0 \rangle, \langle 0, 0, c \rangle \rangle$  holds (the homography of  $N$ )( $P$ ) = Dir111. The theorem is a consequence of (27) and (28).
- (30) Let us consider points  $P_1, P_2, P_3, P_4$  of the projective space over  $\mathcal{E}_T^3$ . Suppose  $P_1, P_2$  and  $P_3$  are not collinear and  $P_1, P_2$  and  $P_4$  are not collinear and  $P_1, P_3$  and  $P_4$  are not collinear and  $P_2, P_3$  and  $P_4$  are not collinear. Then there exists an invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3 such that
- (i) (the homography of  $N$ )( $P_1$ ) = Dir100, and
  - (ii) (the homography of  $N$ )( $P_2$ ) = Dir010, and
  - (iii) (the homography of  $N$ )( $P_3$ ) = Dir001, and
  - (iv) (the homography of  $N$ )( $P_4$ ) = Dir111.

The theorem is a consequence of (24), (29), (9), (25), and (13).

- (31) Let us consider points  $P_1, P_2, P_3, P_4, Q_1, Q_2, Q_3, Q_4$  of the projective space over  $\mathcal{E}_T^3$ . Suppose  $P_1, P_2$  and  $P_3$  are not collinear and  $P_1, P_2$  and  $P_4$  are not collinear and  $P_1, P_3$  and  $P_4$  are not collinear and  $P_2, P_3$  and  $P_4$

are not collinear and  $Q_1$ ,  $Q_2$  and  $Q_3$  are not collinear and  $Q_1$ ,  $Q_2$  and  $Q_4$  are not collinear and  $Q_1$ ,  $Q_3$  and  $Q_4$  are not collinear and  $Q_2$ ,  $Q_3$  and  $Q_4$  are not collinear. Then there exists an invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3 such that

- (i) (the homography of  $N$ )( $P_1$ ) =  $Q_1$ , and
- (ii) (the homography of  $N$ )( $P_2$ ) =  $Q_2$ , and
- (iii) (the homography of  $N$ )( $P_3$ ) =  $Q_3$ , and
- (iv) (the homography of  $N$ )( $P_4$ ) =  $Q_4$ .

The theorem is a consequence of (30), (15), and (13).

#### REFERENCES

- [1] Grzegorz Bancerek and Krzysztof Hryniewiecki. Segments of natural numbers and finite sequences. *Formalized Mathematics*, 1(1):107–114, 1990.
- [2] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [3] Karol Borsuk and Wanda Szmielew. *Foundations of Geometry*. North Holland, 1960.
- [4] Czesław Byliński. The sum and product of finite sequences of real numbers. *Formalized Mathematics*, 1(4):661–668, 1990.
- [5] Roland Coghetto. Homography in  $\mathbb{R}P^2$ . *Formalized Mathematics*, 24(4):239–251, 2016. doi:10.1515/forma-2016-0020.
- [6] Agata Darmochwał. The Euclidean space. *Formalized Mathematics*, 2(4):599–603, 1991.
- [7] Kanchun, Hiroshi Yamazaki, and Yatsuka Nakamura. Cross products and tripple vector products in 3-dimensional Euclidean space. *Formalized Mathematics*, 11(4):381–383, 2003.
- [8] Wojciech Leończuk and Krzysztof Prażmowski. A construction of analytical projective space. *Formalized Mathematics*, 1(4):761–766, 1990.
- [9] Timothy James McKenzie Makarios. The independence of Tarski’s Euclidean Axiom. *Archive of Formal Proofs*, October 2012. Formal proof development.
- [10] Timothy James McKenzie Makarios. A mechanical verification of the independence of Tarski’s Euclidean Axiom. Victoria University of Wellington, New Zealand, 2012. Master’s thesis.
- [11] Jürgen Richter-Gebert. *Perspectives on projective geometry: a guided tour through real and complex geometry*. Springer Science & Business Media, 2011.

Received March 17, 2017



The English version of this volume of Formalized Mathematics was financed under agreement 548/P-DUN/2016 with the funds from the Polish Minister of Science and Higher Education for the dissemination of science.