

# Klein-Beltrami model. Part III

Roland Coghetto   
Rue de la Brasserie 5  
7100 La Louvière, Belgium

**Summary.** Timothy Makarios (with Isabelle/HOL<sup>1</sup>) and John Harrison (with HOL-Light<sup>2</sup>) shown that “the Klein-Beltrami model of the hyperbolic plane satisfy all of Tarski’s axioms except his Euclidean axiom” [2],[3],[4],[5].

With the Mizar system [1] we use some ideas taken from Tim Makarios’s MSc thesis [10] to formalize some definitions (like the absolute) and lemmas necessary for the verification of the independence of the parallel postulate. In this article we prove that our constructed model (we prefer “Beltrami-Klein” name over “Klein-Beltrami”, which can be seen in the naming convention for Mizar functors, and even MML identifiers) satisfies the congruence symmetry, the congruence equivalence relation, and the congruence identity axioms formulated by Tarski (and formalized in Mizar as described briefly in [8]).

MSC: 51A05 51M10 68V20

Keywords: Tarski’s geometry axioms; foundations of geometry; Klein-Beltrami model

MML identifier: BKMODEL3, version: 8.1.09 5.60.1371

## 1. PRELIMINARIES

Now we state the propositions:

- (1) Let us consider real numbers  $x, y$ . If  $x \cdot y < 0$ , then  $0 < \frac{x}{x-y} < 1$ .
- (2) Let us consider a non zero real number  $a$ , and real numbers  $b, r$ . Suppose  $r = \sqrt{a^2 + b^2}$ . Then
  - (i)  $r$  is positive, and

---

<sup>1</sup>[https://www.isa-afp.org/entries/Tarskis\\_Geometry.html](https://www.isa-afp.org/entries/Tarskis_Geometry.html)

<sup>2</sup><https://github.com/jrh13/hol-light/blob/master/100/independence.ml>

- (ii)  $(\frac{a}{r})^2 + (\frac{b}{r})^2 = 1$ .
- (3) Let us consider a non zero real number  $a$ , and real numbers  $b, c, d, e$ .  
Suppose  $a \cdot b = c - d \cdot e$ . Then  $b^2 = \frac{c^2}{a^2} - 2 \cdot \frac{c \cdot d}{a \cdot a} \cdot e + \frac{d^2}{a^2} \cdot e^2$ .  
Let us consider complex numbers  $a, b, c$ . Now we state the propositions:
- (4) If  $a \neq 0$ , then  $\frac{a^2 \cdot b \cdot c}{a^2} = b \cdot c$ .
- (5) If  $a \neq 0$ , then  $\frac{2 \cdot a^2 \cdot b \cdot c}{a^2} = 2 \cdot b \cdot c$ . The theorem is a consequence of (4).
- (6) Let us consider a real number  $a$ . If  $1 < a$ , then  $\frac{1}{a} - 1 < 0$ .
- (7) Let us consider real numbers  $a, b$ . If  $0 < a$  and  $1 < b$ , then  $\frac{a}{b} - a < 0$ .  
The theorem is a consequence of (6).
- (8) Let us consider a non zero real number  $a$ , and real numbers  $b, c, d$ .  
Suppose  $a^2 + c^2 = b^2$  and  $1 < b^2$ . Then  $\frac{(b^2)^2}{a^2} - 2 \cdot \frac{b^2 \cdot c}{a \cdot a} \cdot d + \frac{c^2}{a^2} \cdot d^2 + d^2 \neq 1$ .  
The theorem is a consequence of (5) and (7).
- (9) Let us consider real numbers  $a, b, c$ . If  $a \cdot (-b) = c$  and  $a \cdot c = b$ , then  $c = 0$  and  $b = 0$ .
- (10) Let us consider a positive real number  $a$ . Then  $\frac{\sqrt{a}}{a} = \frac{1}{\sqrt{a}}$ .

## 2. PLANAR LEMMAS

Let  $a$  be a non zero real number and  $b, c$  be real numbers. Observe that  $[a, b, c]$  is non zero as an element of  $\mathcal{E}_T^3$  and  $[c, a, b]$  is non zero as an element of  $\mathcal{E}_T^3$  and  $[b, c, a]$  is non zero as an element of  $\mathcal{E}_T^3$ .

Let  $P$  be an element of the real projective plane. Assume  $P \in (\text{the absolute}) \cup (\text{the BK-model})$ . The functor  $\#P$  yielding a non zero element of  $\mathcal{E}_T^3$  is defined by

(Def. 1) the direction of  $it = P$  and  $it(3) = 1$ .

Now we state the propositions:

- (11) Let us consider an element  $P$  of the real projective plane. Then there exists an element  $Q$  of the BK-model such that  $P \neq Q$ .

From now on  $P$  denotes an element of the BK-model.

- (12) There exist elements  $P_1, P_2$  of the absolute such that

- (i)  $P_1 \neq P_2$ , and  
(ii)  $P_1, P$  and  $P_2$  are collinear.

The theorem is a consequence of (11).

- (13) Let us consider an element  $Q$  of the absolute. Then there exists an element  $R$  of the BK-model such that

- (i)  $P \neq R$ , and

(ii)  $P, Q$  and  $R$  are collinear.

(14) Let us consider a line  $L$  of Inc- $\text{ProjSp}$ (the real projective plane). Suppose  $P \in L$ . Then there exist elements  $P_1, P_2$  of the absolute such that

(i)  $P_1 \neq P_2$ , and

(ii)  $P_1, P_2 \in L$ .

Let  $N$  be an invertible square matrix over  $\mathbb{R}_F$  of dimension 3. The functor Line-homography( $N$ ) yielding a function from the lines of Inc- $\text{ProjSp}$ (the real projective plane) into the lines of Inc- $\text{ProjSp}$ (the real projective plane) is defined by

(Def. 2) for every line  $x$  of Inc- $\text{ProjSp}$ (the real projective plane),  $it(x) = \{(the\ homography\ of\ N)(P),\ where\ P\ is\ a\ point\ of\ Inc-ProjSp(the\ real\ projective\ plane) : P\ lies\ on\ x\}$ .

In the sequel  $N, N_1, N_2$  denote invertible square matrices over  $\mathbb{R}_F$  of dimension 3 and  $l, l_1, l_2$  denote elements of the lines of Inc- $\text{ProjSp}$ (the real projective plane). Now we state the propositions:

(15)  $(\text{Line-homography}(N_1))((\text{Line-homography}(N_2))(l)) = (\text{Line-homography}(N_1 \cdot N_2))(l)$ .

PROOF: Reconsider  $l_2 = (\text{Line-homography}(N_2))(l)$  as a line of Inc- $\text{ProjSp}$ (the real projective plane).  $\{(the\ homography\ of\ N_1)(P),\ where\ P\ is\ a\ point\ of\ Inc-ProjSp(the\ real\ projective\ plane) : P\ lies\ on\ l_2\} = \{(the\ homography\ of\ N_1 \cdot N_2)(P),\ where\ P\ is\ a\ point\ of\ Inc-ProjSp(the\ real\ projective\ plane) : P\ lies\ on\ l\}$  by [9, (3), (4), (5)], [6, (13)].  $\square$

(16)  $(\text{Line-homography}(I_{\mathbb{R}_F}^{3 \times 3}))(l) = l$ .

PROOF: Set  $X = \{(the\ homography\ of\ I_{\mathbb{R}_F}^{3 \times 3})(P),\ where\ P\ is\ a\ point\ of\ Inc-ProjSp(the\ real\ projective\ plane) : P\ lies\ on\ l\}$ .  $X \subseteq l, l \subseteq X$ .  $\square$

(17) (i)  $(\text{Line-homography}(N))((\text{Line-homography}(N^\sim))(l)) = l$ , and

(ii)  $(\text{Line-homography}(N^\sim))((\text{Line-homography}(N))(l)) = l$ .

The theorem is a consequence of (15) and (16).

(18) If  $(\text{Line-homography}(N))(l_1) = (\text{Line-homography}(N))(l_2)$ , then  $l_1 = l_2$ .

The theorem is a consequence of (17).

The functor SetLineHom3 yielding a set is defined by the term

(Def. 3) the set of all Line-homography( $N$ ) where  $N$  is an invertible square matrix over  $\mathbb{R}_F$  of dimension 3.

Observe that SetLineHom3 is non empty. Let  $h_1, h_2$  be elements of SetLineHom3. The functor  $h_1 \circ h_2$  yielding an element of SetLineHom3 is defined by

(Def. 4) there exist invertible square matrices  $N_1, N_2$  over  $\mathbb{R}_F$  of dimension 3 such that  $h_1 = \text{Line-homography}(N_1)$  and  $h_2 = \text{Line-homography}(N_2)$  and  $it = \text{Line-homography}(N_1 \cdot N_2)$ .

Now we state the propositions:

(19) Let us consider elements  $h_1, h_2$  of  $\text{SetLineHom3}$ . Suppose  $h_1 = \text{Line-homography}(N_1)$  and  $h_2 = \text{Line-homography}(N_2)$ .

Then  $\text{Line-homography}(N_1 \cdot N_2) = h_1 \circ h_2$ . The theorem is a consequence of (15).

(20) Let us consider elements  $x, y, z$  of  $\text{SetLineHom3}$ . Then  $(x \circ y) \circ z = x \circ (y \circ z)$ . The theorem is a consequence of (19).

The functor  $\text{BinOpLineHom3}$  yielding a binary operation on  $\text{SetLineHom3}$  is defined by

(Def. 5) for every elements  $h_1, h_2$  of  $\text{SetLineHom3}$ ,  $it(h_1, h_2) = h_1 \circ h_2$ .

The functor  $\text{GroupLineHom3}$  yielding a strict multiplicative magma is defined by the term

(Def. 6)  $\langle \text{SetLineHom3}, \text{BinOpLineHom3} \rangle$ .

Let us observe that  $\text{GroupLineHom3}$  is non empty, associative, and group-like. Now we state the propositions:

(21)  $\mathbf{1}_{\text{GroupLineHom3}} = \text{Line-homography}(I_{\mathbb{R}_F}^{3 \times 3})$ .

(22) Let us consider elements  $h, g$  of  $\text{GroupLineHom3}$ , and invertible square matrices  $N, N_1$  over  $\mathbb{R}_F$  of dimension 3. Suppose  $h = \text{Line-homography}(N)$  and  $g = \text{Line-homography}(N_1)$  and  $N_1 = N^\sim$ . Then  $g = h^{-1}$ . The theorem is a consequence of (21).

In the sequel  $P$  denotes a point of the projective space over  $\mathcal{E}_T^3$  and  $l$  denotes a line of  $\text{Inc-ProjSp}$ (the real projective plane).

(23) If (the homography of  $N$ )( $P$ )  $\in l$ , then  $P \in (\text{Line-homography}(N^\sim))(l)$ .

(24) If  $P \in (\text{Line-homography}(N))(l)$ , then (the homography of  $N^\sim$ )( $P$ )  $\in l$ .

(25)  $P \in l$  if and only if (the homography of  $N$ )( $P$ )  $\in (\text{Line-homography}(N))(l)$ . The theorem is a consequence of (23) and (17).

(26) Let us consider non zero elements  $u, v, w$  of  $\mathcal{E}_T^3$ . Suppose  $(u)_3 = 1$  and  $(v)_1 = -(u)_2$  and  $(v)_2 = (u)_1$  and  $(v)_3 = 0$  and  $(w)_3 = 1$  and  $\langle |u, v, w| \rangle = 0$ . Then  $((u)_1)^2 + ((u)_2)^2 - (u)_1 \cdot (w)_1 - (u)_2 \cdot (w)_2 = 0$ .

(27) Let us consider a non zero real number  $a$ , and real numbers  $b, c$ . Then  $a \cdot [\frac{b}{a}, \frac{c}{a}, 1] = [b, c, a]$ .

Let us consider non zero elements  $u, v, w$  of  $\mathcal{E}_T^3$ . Now we state the propositions:

- (28) Suppose  $(u)_1 \neq 0$  and  $(u)_3 = 1$  and  $(v)_1 = -(u)_2$  and  $(v)_2 = (u)_1$  and  $(v)_3 = 0$  and  $(w)_3 = 1$  and  $\langle |u, v, w| \rangle = 0$  and  $1 < ((u)_1)^2 + ((u)_2)^2$ . Then  $((w)_1)^2 + ((w)_2)^2 \neq 1$ . The theorem is a consequence of (26), (2), (3), and (8).
- (29) Suppose  $(u)_2 \neq 0$  and  $(u)_3 = 1$  and  $(v)_1 = -(u)_2$  and  $(v)_2 = (u)_1$  and  $(v)_3 = 0$  and  $(w)_3 = 1$  and  $\langle |u, v, w| \rangle = 0$  and  $1 < ((u)_1)^2 + ((u)_2)^2$ . Then  $((w)_1)^2 + ((w)_2)^2 \neq 1$ . The theorem is a consequence of (26), (2), (3), and (8).
- (30) Let us consider an element  $P$  of the absolute. Then there exists a non zero element  $u$  of  $\mathcal{E}_T^3$  such that
- (i)  $u(3) = 1$ , and
  - (ii)  $P =$  the direction of  $u$ .
- (31) Let us consider real numbers  $a, b, c, d$ , and non zero elements  $u, v$  of  $\mathcal{E}_T^3$ . Suppose  $u = [a, b, 1]$  and  $v = [c, d, 0]$ . Then the direction of  $u \neq$  the direction of  $v$ .
- (32) Let us consider a non zero element  $u$  of  $\mathcal{E}_T^3$ . Suppose  $u(1)^2 + u(2)^2 < 1$  and  $u(3) = 1$ . Then the direction of  $u$  is an element of the BK-model.
- (33) Let us consider real numbers  $a, b$ . Suppose  $a^2 + b^2 \leq 1$ . Then the direction of  $[a, b, 1] \in$  (the BK-model)  $\cup$  (the absolute). The theorem is a consequence of (32).
- (34) If  $P \notin$  (the BK-model)  $\cup$  (the absolute), then there exists  $l$  such that  $P \in l$  and  $l$  misses the absolute. The theorem is a consequence of (9), (30), (27), (31), (33), (28), and (29).
- (35) Let us consider a point  $P$  of the real projective plane, an element  $h$  of the subgroup of  $K$ -isometries, and an invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3. Suppose  $h =$  the homography of  $N$ . Then  $P$  is an element of the absolute if and only if (the homography of  $N$ )( $P$ ) is an element of the absolute.

Let us consider an element  $P$  of the BK-model, an element  $h$  of the subgroup of  $K$ -isometries, and an invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3.

- (36) If  $h =$  the homography of  $N$ , then (the homography of  $N$ )( $P$ ) is an element of the BK-model.

PROOF: Set  $h_1 =$  (the homography of  $N$ )( $P$ ).  $h_1$  is not an element of the absolute by (35), [7, (1)]. Consider  $l$  such that  $h_1 \in l$  and  $l$  misses the absolute. Reconsider  $L =$  (Line-homography( $N^\sim$ ))( $l$ ) as a line of the real projective plane. Reconsider  $L' = L$  as a line of Inc-ProjSp(the real projective plane). Consider  $P_1, P_2$  being elements of the absolute such that

$P_1 \neq P_2$  and  $P_1 \in L'$  and  $P_2 \in L'$ . (The homography of  $N$ )( $P_1$ ) is an element of the absolute. (The homography of  $N$ )( $P_1$ )  $\in$  (Line-homography( $N$ ))( $L$ ). (The homography of  $N$ )( $P_1$ )  $\in$   $l$ .  $\square$

(37) Suppose  $h$  = the homography of  $N$ . Then there exists a non zero element  $u$  of  $\mathcal{E}_T^3$  such that

- (i) (the homography of  $N$ )( $P$ ) = the direction of  $u$ , and
- (ii)  $u(3) = 1$ .

The theorem is a consequence of (36).

### 3. THE CONSTRUCTION OF BELTRAMI-KLEIN MODEL

The functor BK-model-Betweenness yielding a relation between (the BK-model)  $\times$  (the BK-model) and the BK-model is defined by

(Def. 7) for every elements  $a, b, c$  of the BK-model,  $\langle\langle a, b \rangle, c \rangle \in it$  iff  $\text{BK-to-REAL2}(b) \in \mathcal{L}(\text{BK-to-REAL2}(a), \text{BK-to-REAL2}(c))$ .

The functor BK-model-Equidistance yielding a relation between (the BK-model)  $\times$  (the BK-model) and (the BK-model)  $\times$  (the BK-model) is defined by

(Def. 8) for every elements  $a, b, c, d$  of the BK-model,  $\langle\langle a, b \rangle, \langle c, d \rangle \rangle \in it$  iff there exists an element  $h$  of the subgroup of  $K$ -isometries and there exists an invertible square matrix  $N$  over  $\mathbb{R}_F$  of dimension 3 such that  $h$  = the homography of  $N$  and (the homography of  $N$ )( $a$ ) =  $c$  and (the homography of  $N$ )( $b$ ) =  $d$ .

The functor BK-model-Plane yielding a Tarski plane is defined by the term

(Def. 9)  $\langle$ the BK-model, BK-model-Betweenness, BK-model-Equidistance $\rangle$ .

### 4. CONGRUENCE SYMMETRY

Now we state the proposition:

(38) BK-model-Plane satisfies the axiom of congruence symmetry.

### 5. CONGRUENCE EQUIVALENCE RELATION

Now we state the proposition:

(39) BK-model-Plane satisfies the axiom of congruence equivalence relation.

## 6. CONGRUENCE IDENTITY

Now we state the proposition:

- (40) BK-model-Plane satisfies the axiom of congruence identity.

## REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Kornilowicz, Roman Matuszewski, Adam Naumowicz, Karol Pał, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8\_17.
- [2] Eugenio Beltrami. Saggio di interpretazione della geometria non-euclidea. *Giornale di Matematiche*, 6:284–322, 1868.
- [3] Eugenio Beltrami. Essai d'interprétation de la géométrie non-euclidéenne. In *Annales scientifiques de l'École Normale Supérieure. Trad. par J. Houël*, volume 6, pages 251–288. Elsevier, 1869.
- [4] Karol Borsuk and Wanda Szmielew. *Foundations of Geometry*. North Holland, 1960.
- [5] Karol Borsuk and Wanda Szmielew. *Podstawy geometrii*. Państwowe Wydawnictwo Naukowe, Warszawa, 1955 (in Polish).
- [6] Roland Coghetto. Group of homography in real projective plane. *Formalized Mathematics*, 25(1):55–62, 2017. doi:10.1515/forma-2017-0005.
- [7] Roland Coghetto. Klein-Beltrami model. Part II. *Formalized Mathematics*, 26(1):33–48, 2018. doi:10.2478/forma-2018-0004.
- [8] Adam Grabowski and Roland Coghetto. Tarski's geometry and the Euclidean plane in Mizar. In *Joint Proceedings of the FM4M, MathUI, and ThEdu Workshops, Doctoral Program, and Work in Progress at the Conference on Intelligent Computer Mathematics 2016 co-located with the 9th Conference on Intelligent Computer Mathematics (CICM 2016), Białystok, Poland, July 25–29, 2016*, volume 1785 of *CEUR-WS*, pages 4–9. CEUR-WS.org, 2016.
- [9] Wojciech Leńczuk and Krzysztof Prażmowski. Incidence projective spaces. *Formalized Mathematics*, 2(2):225–232, 1991.
- [10] Timothy James McKenzie Makarios. A mechanical verification of the independence of Tarski's Euclidean Axiom. Victoria University of Wellington, New Zealand, 2012. Master's thesis.

*Accepted December 30, 2019*

---