

Klein-Beltrami model. Part IV

Roland Coghetto 
Rue de la Brasserie 5
7100 La Louvière, Belgium

Summary. Timothy Makarios (with Isabelle/HOL¹) and John Harrison (with HOL-Light²) shown that “the Klein-Beltrami model of the hyperbolic plane satisfy all of Tarski’s axioms except his Euclidean axiom” [2],[3],[4, 5].

With the Mizar system [1] we use some ideas taken from Tim Makarios’s MSc thesis [10] to formalize some definitions and lemmas necessary for the verification of the independence of the parallel postulate. In this article, which is the continuation of [8], we prove that our constructed model satisfies the axioms of segment construction, the axiom of betweenness identity, and the axiom of Pasch due to Tarski, as formalized in [11] and related Mizar articles.

MSC: 51A05 51M10 68V20

Keywords: Tarski’s geometry axioms; foundations of geometry; Klein-Beltrami model

MML identifier: BKMODEL4, version: 8.1.09 5.60.1371

1. PRELIMINARIES

Let us consider real numbers a, b . Now we state the propositions:

- (1) If $a \neq b$, then $1 - \frac{a}{a-b} = -\frac{b}{a-b}$.
- (2) If $0 < a \cdot b$, then $0 < \frac{a}{b}$.

Now we state the propositions:

- (3) Let us consider real numbers a, b, c . Suppose $0 \leq a \leq 1$ and $0 < b \cdot c$. Then $0 \leq \frac{a \cdot c}{(1-a) \cdot b + a \cdot c} \leq 1$.

¹https://www.isa-afp.org/entries/Tarskis_Geometry.html

²<https://github.com/jrh13/hol-light/blob/master/100/independence.ml>

- (4) Let us consider real numbers a, b, c . Suppose $(1-a) \cdot b + a \cdot c \neq 0$. Then $1 - \frac{a \cdot c}{(1-a) \cdot b + a \cdot c} = \frac{(1-a) \cdot b}{(1-a) \cdot b + a \cdot c}$.
- (5) Let us consider real numbers a, b, c, d . If $b \neq 0$, then $\frac{a \cdot b \cdot d}{b} = \frac{a \cdot d}{c}$.
- (6) Let us consider an element u of \mathcal{E}_T^3 . Then $u = [u(1), u(2), u(3)]$.
- (7) Let us consider an element P of the BK-model. Then $\text{BK-to-REAL2}(P) \in \text{TarskiEuclid2Space}$.

Let P be a point of BK-model-Plane. The functor $\text{BKtoT2}(P)$ yielding a point of $\text{TarskiEuclid2Space}$ is defined by

- (Def. 1) there exists an element p of the BK-model such that $P = p$ and $it = \text{BK-to-REAL2}(p)$.

Let P be a point of $\text{TarskiEuclid2Space}$. Assume $\hat{P} \in$ the inside of $\text{circle}(0,0,1)$. The functor $\text{T2toBK}(P)$ yielding a point of BK-model-Plane is defined by

- (Def. 2) there exists a non zero element u of \mathcal{E}_T^3 such that $it =$ the direction of u and $(u)_3 = 1$ and $\hat{P} = [(u)_1, (u)_2]$.

Let us consider a point P of BK-model-Plane. Now we state the propositions:

- (8) $\text{BKto}\hat{\text{T2}}(P) \in$ the inside of $\text{circle}(0,0,1)$.
- (9) $\text{T2toBK}(\text{BKtoT2}(P)) = P$.
- (10) Let us consider a point P of $\text{TarskiEuclid2Space}$. Suppose \hat{P} is an element of the inside of $\text{circle}(0,0,1)$. Then $\text{BKtoT2}(\text{T2toBK}(P)) = P$.
- (11) Let us consider a point P of BK-model-Plane, and an element p of the BK-model. Suppose $P = p$. Then
- (i) $\text{BKtoT2}(P) = \text{BK-to-REAL2}(p)$, and
 - (ii) $\text{BKto}\hat{\text{T2}}(P) = \text{BK-to-REAL2}(p)$.
- (12) Let us consider points P, Q, R of BK-model-Plane, and points P_2, Q_2, R_2 of $\text{TarskiEuclid2Space}$. Suppose $P_2 = \text{BKtoT2}(P)$ and $Q_2 = \text{BKtoT2}(Q)$ and $R_2 = \text{BKtoT2}(R)$. Then Q lies between P and R if and only if Q_2 lies between P_2 and R_2 . The theorem is a consequence of (11).
- (13) Let us consider elements P, Q of \mathcal{E}_T^2 . If $P \neq Q$, then $P(1) \neq Q(1)$ or $P(2) \neq Q(2)$.
- (14) Let us consider real numbers a, b , and elements P, Q of \mathcal{E}_T^2 . If $P \neq Q$ and $(1-a) \cdot P + a \cdot Q = (1-b) \cdot P + b \cdot Q$, then $a = b$. The theorem is a consequence of (13).
- (15) Let us consider points P, Q of BK-model-Plane. If $\text{BKto}\hat{\text{T2}}(P) = \text{BKto}\hat{\text{T2}}(Q)$, then $P = Q$. The theorem is a consequence of (11).

Let P, Q, R be points of BK-model-Plane. Assume Q lies between P and R and $P \neq R$. The functor $\text{length}(P, Q, R)$ yielding a real number is defined by

(Def. 3) $0 \leq it \leq 1$ and $\text{BKto}\hat{\text{T}}2(Q) = (1 - it) \cdot (\text{BKto}\hat{\text{T}}2(P)) + it \cdot (\text{BKto}\hat{\text{T}}2(R))$.

Let us consider points P, Q of BK-model-Plane. Now we state the propositions:

(16) (i) P lies between P and Q , and

(ii) Q lies between P and Q .

The theorem is a consequence of (12).

(17) If $P \neq Q$, then $\text{length}(P, P, Q) = 0$ and $\text{length}(P, Q, Q) = 1$. The theorem is a consequence of (16).

(18) Let us consider a square matrix N over \mathbb{R}_F of dimension 3. Suppose $N = \langle \langle 2, 0, -1 \rangle, \langle 0, \sqrt{3}, 0 \rangle, \langle 1, 0, -2 \rangle \rangle$. Then

(i) $\text{Det } N = (-3) \cdot \sqrt{3}$, and

(ii) N is invertible.

(19) Let us consider elements $a_{11}, a_{12}, a_{13}, a_{21}, a_{22}, a_{23}, a_{31}, a_{32}, a_{33}, b_{11}, b_{12}, b_{13}, b_{21}, b_{22}, b_{23}, b_{31}, b_{32}, b_{33}, a_1, a_2, a_3, a_4, a_5, a_6, a_7, a_8, a_9$ of \mathbb{R}_F , and square matrices A, B over \mathbb{R}_F of dimension 3.

Suppose $A = \langle \langle a_{11}, a_{12}, a_{13} \rangle, \langle a_{21}, a_{22}, a_{23} \rangle, \langle a_{31}, a_{32}, a_{33} \rangle \rangle$ and $B = \langle \langle b_{11}, b_{12}, b_{13} \rangle, \langle b_{21}, b_{22}, b_{23} \rangle, \langle b_{31}, b_{32}, b_{33} \rangle \rangle$ and $a_1 = a_{11} \cdot b_{11} + a_{12} \cdot b_{21} + a_{13} \cdot b_{31}$ and $a_2 = a_{11} \cdot b_{12} + a_{12} \cdot b_{22} + a_{13} \cdot b_{32}$ and $a_3 = a_{11} \cdot b_{13} + a_{12} \cdot b_{23} + a_{13} \cdot b_{33}$ and $a_4 = a_{21} \cdot b_{11} + a_{22} \cdot b_{21} + a_{23} \cdot b_{31}$.

Suppose $a_5 = a_{21} \cdot b_{12} + a_{22} \cdot b_{22} + a_{23} \cdot b_{32}$ and $a_6 = a_{21} \cdot b_{13} + a_{22} \cdot b_{23} + a_{23} \cdot b_{33}$ and $a_7 = a_{31} \cdot b_{11} + a_{32} \cdot b_{21} + a_{33} \cdot b_{31}$ and $a_8 = a_{31} \cdot b_{12} + a_{32} \cdot b_{22} + a_{33} \cdot b_{32}$ and $a_9 = a_{31} \cdot b_{13} + a_{32} \cdot b_{23} + a_{33} \cdot b_{33}$.

Then $A \cdot B = \langle \langle a_1, a_2, a_3 \rangle, \langle a_4, a_5, a_6 \rangle, \langle a_7, a_8, a_9 \rangle \rangle$.

Let us consider square matrices N_1, N_2 over \mathbb{R}_F of dimension 3. Now we state the propositions:

(20) Suppose $N_1 = \langle \langle 2, 0, -1 \rangle, \langle 0, \sqrt{3}, 0 \rangle, \langle 1, 0, -2 \rangle \rangle$ and $N_2 = \langle \langle \frac{2}{3}, 0, -\frac{1}{3} \rangle, \langle 0, \frac{1}{\sqrt{3}}, 0 \rangle, \langle \frac{1}{3}, 0, -\frac{2}{3} \rangle \rangle$. Then $N_1 \cdot N_2 = \langle \langle 1, 0, 0 \rangle, \langle 0, 1, 0 \rangle, \langle 0, 0, 1 \rangle \rangle$. The theorem is a consequence of (19).

(21) Suppose $N_2 = \langle \langle 2, 0, -1 \rangle, \langle 0, \sqrt{3}, 0 \rangle, \langle 1, 0, -2 \rangle \rangle$ and $N_1 = \langle \langle \frac{2}{3}, 0, -\frac{1}{3} \rangle, \langle 0, \frac{1}{\sqrt{3}}, 0 \rangle, \langle \frac{1}{3}, 0, -\frac{2}{3} \rangle \rangle$. Then $N_1 \cdot N_2 = \langle \langle 1, 0, 0 \rangle, \langle 0, 1, 0 \rangle, \langle 0, 0, 1 \rangle \rangle$. The theorem is a consequence of (19).

(22) Suppose $N_1 = \langle \langle 2, 0, -1 \rangle, \langle 0, \sqrt{3}, 0 \rangle, \langle 1, 0, -2 \rangle \rangle$ and $N_2 = \langle \langle \frac{2}{3}, 0, -\frac{1}{3} \rangle, \langle 0, \frac{1}{\sqrt{3}}, 0 \rangle, \langle \frac{1}{3}, 0, -\frac{2}{3} \rangle \rangle$. Then N_1 is inverse of N_2 . The theorem is a consequence of (20) and (21).

Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3. Now we state the propositions:

- (23) Suppose $N = \langle \langle \frac{2}{3}, 0, -\frac{1}{3} \rangle, \langle 0, \frac{1}{\sqrt{3}}, 0 \rangle, \langle \frac{1}{3}, 0, -\frac{2}{3} \rangle \rangle$. Then (the homography of N) $^\circ$ (the absolute) \subseteq the absolute.
 PROOF: (The homography of N) $^\circ$ (the absolute) \subseteq the absolute by [7, (89)], [9, (7)]. \square
- (24) Suppose $N = \langle \langle 2, 0, -1 \rangle, \langle 0, \sqrt{3}, 0 \rangle, \langle 1, 0, -2 \rangle \rangle$. Then (the homography of N) $^\circ$ (the absolute) = the absolute.
 PROOF: (The homography of N) $^\circ$ (the absolute) \subseteq the absolute.
 The absolute \subseteq (the homography of N) $^\circ$ (the absolute) by [6, (19)], (22), (23). \square
- (25) Let us consider real numbers a, b, r , and elements P, Q, R of \mathcal{E}_T^2 . Suppose $Q \in \mathcal{L}(P, R)$ and $P, R \in$ the inside of circle(a, b, r). Then $Q \in$ the inside of circle(a, b, r).
- (26) Let us consider non zero elements u, v of \mathcal{E}_T^3 . Suppose the direction of $u =$ the direction of v and $u(3) \neq 0$ and $u(3) = v(3)$. Then $u = v$.
- (27) Let us consider an element R of the projective space over \mathcal{E}_T^3 , elements P, Q of the BK-model, non zero elements u, v, w of \mathcal{E}_T^3 , and a real number r . Suppose $0 \leq r \leq 1$ and $P =$ the direction of u and $Q =$ the direction of v and $R =$ the direction of w and $u(3) = 1$ and $v(3) = 1$ and $w = r \cdot u + (1 - r) \cdot v$. Then R is an element of the BK-model.
 PROOF: Consider u_2 being a non zero element of \mathcal{E}_T^3 such that the direction of $u_2 = P$ and $u_2(3) = 1$ and BK-to-REAL2(P) = $[u_2(1), u_2(2)]$. $u = u_2$. Reconsider $r_4 = [u_2(1), u_2(2)]$ as an element of \mathcal{E}_T^2 . Consider v_2 being a non zero element of \mathcal{E}_T^3 such that the direction of $v_2 = Q$ and $v_2(3) = 1$ and BK-to-REAL2(Q) = $[v_2(1), v_2(2)]$. $v = v_2$. Reconsider $r_6 = [v_2(1), v_2(2)]$ as an element of \mathcal{E}_T^2 . Reconsider $r_8 = [w(1), w(2)]$ as an element of \mathcal{E}_T^2 . $r_8 = r \cdot r_4 + (1 - r) \cdot r_6$. Consider R_3 being an element of \mathcal{E}_T^2 such that $R_3 = r_8$ and REAL2-to-BK(r_8) = the direction of $[(R_3)_1, (R_3)_2, 1]$. \square
- (28) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , points P, Q of the projective space over \mathcal{E}_T^3 , and non zero elements u, v of \mathcal{E}_T^3 . Suppose $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $P =$ the direction of u and $Q =$ the direction of v and $Q =$ (the homography of N)(P) and $u(3) = 1$. Then there exists a non zero real number a such that
- (i) $v(1) = a \cdot (n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13})$, and
 - (ii) $v(2) = a \cdot (n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23})$, and
 - (iii) $v(3) = a \cdot (n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33})$.
- (29) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , an element

P of the BK-model, a point Q of the projective space over \mathcal{E}_T^3 , and non zero elements u, v of \mathcal{E}_T^3 . Suppose $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $P =$ the direction of u and $Q =$ the direction of v and $Q =$ (the homography of N)(P) and $u(3) = 1$ and $v(3) = 1$. Then

- (i) $n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33} \neq 0$, and
- (ii) $v(1) = \frac{n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}$, and
- (iii) $v(2) = \frac{n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}$.

The theorem is a consequence of (28).

- (30) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, an element h of the subgroup of K -isometries, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , an element P of the BK-model, and a non zero element u of \mathcal{E}_T^3 . Suppose $h =$ the homography of N and $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $P =$ the direction of u and $u(3) = 1$. Then $n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33} \neq 0$. The theorem is a consequence of (29).

- (31) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , an element P of the absolute, a point Q of the projective space over \mathcal{E}_T^3 , and non zero elements u, v of \mathcal{E}_T^3 . Suppose $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $P =$ the direction of u and $Q =$ the direction of v and $Q =$ (the homography of N)(P) and $u(3) = 1$ and $v(3) = 1$. Then

- (i) $n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33} \neq 0$, and
- (ii) $v(1) = \frac{n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}$, and
- (iii) $v(2) = \frac{n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}$.

The theorem is a consequence of (28).

- (32) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, an element h of the subgroup of K -isometries, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , an element P of the absolute, and a non zero element u of \mathcal{E}_T^3 . Suppose $h =$ the homography of N and $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $P =$ the direction of u and $u(3) = 1$. Then $n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33} \neq 0$. The theorem is a consequence of (31).

- (33) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, an element h of the subgroup of K -isometries, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , an element P of the BK-model, and a non zero element u of \mathcal{E}_T^3 . Suppose $h =$ the homography of N and $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $P =$ the direction

of u and $u(3) = 1$. Then (the homography of N)(P) = the direction of $\left[\frac{n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}, \frac{n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}, 1 \right]$. The theorem is a consequence of (29).

(34) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, an element h of the subgroup of K -isometries, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , an element P of the absolute, and a non zero element u of \mathcal{E}_T^3 . Suppose $h =$ the homography of N and $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $P =$ the direction of u and $u(3) = 1$. Then (the homography of N)(P) = the direction of $\left[\frac{n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}, \frac{n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23}}{n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}}, 1 \right]$. The theorem is a consequence of (31).

(35) Let us consider a subset A of \mathcal{E}_T^3 , a convex, non empty subset B of \mathcal{E}_T^2 , a real number r , and an element x of \mathcal{E}_T^3 . Suppose $A = \{x$, where x is an element of $\mathcal{E}_T^3 : [(x)_1, (x)_2] \in B$ and $(x)_3 = r\}$. Then A is non empty and convex.

(36) Let us consider elements n_1, n_2, n_3 of \mathbb{R}_F , and elements n, u of \mathcal{E}_T^3 . Suppose $n = \langle n_1, n_2, n_3 \rangle$ and $u(3) = 1$. Then $|(n, u)| = n_1 \cdot u(1) + n_2 \cdot u(2) + n_3$.

(37) Let us consider a convex, non empty subset A of \mathcal{E}_T^3 , and elements n, u, v of \mathcal{E}_T^3 . Suppose for every element w of \mathcal{E}_T^3 such that $w \in A$ holds $|(n, w)| \neq 0$ and $u, v \in A$. Then $0 < |(n, u)| \cdot |(n, v)|$.

PROOF: Set $x = |(n, u)|$. Set $y = |(n, v)|$. Reconsider $l = \frac{x}{x-y}$ as a non zero real number. Reconsider $w = l \cdot v + (1-l) \cdot u$ as an element of \mathcal{E}_T^3 . $x \neq y$. $1-l = -\frac{y}{x-y}$. $|(n, w)| = 0$. \square

Let us consider elements n_{31}, n_{32}, n_{33} of \mathbb{R}_F and elements u, v of \mathcal{E}_T^2 . Now we state the propositions:

(38) Suppose $u, v \in$ the inside of circle(0,0,1) and for every element w of \mathcal{E}_T^2 such that $w \in$ the inside of circle(0,0,1) holds $n_{31} \cdot w(1) + n_{32} \cdot w(2) + n_{33} \neq 0$. Then $0 < (n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}) \cdot (n_{31} \cdot v(1) + n_{32} \cdot v(2) + n_{33})$. The theorem is a consequence of (35), (36), and (37).

(39) Suppose $u \in$ the inside of circle(0,0,1) and $v \in$ circle(0,0,1) and for every element w of \mathcal{E}_T^2 such that $w \in$ the closed inside of circle(0,0,1) holds $n_{31} \cdot w(1) + n_{32} \cdot w(2) + n_{33} \neq 0$. Then $0 < (n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}) \cdot (n_{31} \cdot v(1) + n_{32} \cdot v(2) + n_{33})$. The theorem is a consequence of (35), (36), and (37).

(40) Let us consider real numbers l, r , elements u, v, w of \mathcal{E}_T^3 , and real numbers $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}, m_1, m_2, m_3, m_4, m_5, m_6, m_7, m_8, m_9$.

Suppose $m_3 \neq 0$ and $m_6 \neq 0$ and $m_9 \neq 0$ and $r = \frac{l \cdot m_6}{(1-l) \cdot m_3 + l \cdot m_6}$ and $(1-l) \cdot m_3 + l \cdot m_6 \neq 0$ and $w = (1-l) \cdot u + l \cdot v$ and $m_1 = n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13}$ and $m_2 = n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23}$ and $m_3 = n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}$ and $m_4 = n_{11} \cdot v(1) + n_{12} \cdot v(2) + n_{13}$.

Suppose $m_5 = n_{21} \cdot v(1) + n_{22} \cdot v(2) + n_{23}$ and $m_6 = n_{31} \cdot v(1) + n_{32} \cdot v(2) + n_{33}$ and $m_7 = n_{11} \cdot w(1) + n_{12} \cdot w(2) + n_{13}$ and $m_8 = n_{21} \cdot w(1) + n_{22} \cdot w(2) + n_{23}$ and $m_9 = n_{31} \cdot w(1) + n_{32} \cdot w(2) + n_{33}$.

Then $(1-r) \cdot [\frac{m_1}{m_3}, \frac{m_2}{m_3}, 1] + r \cdot [\frac{m_4}{m_6}, \frac{m_5}{m_6}, 1] = [\frac{m_7}{m_9}, \frac{m_8}{m_9}, 1]$. The theorem is a consequence of (4) and (5).

- (41) Let us consider an invertible square matrix N over \mathbb{R}_F of dimension 3, an element h of the subgroup of K -isometries, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , and an element P of the BK-model. Suppose $h =$ the homography of N and $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$. Then (the homography of N)(P) = the direction of $[\frac{n_{11} \cdot (\text{BK-to-REAL2}(P))_1 + n_{12} \cdot (\text{BK-to-REAL2}(P))_2 + n_{13}}{n_{31} \cdot (\text{BK-to-REAL2}(P))_1 + n_{32} \cdot (\text{BK-to-REAL2}(P))_2 + n_{33}}, \frac{n_{21} \cdot (\text{BK-to-REAL2}(P))_1 + n_{22} \cdot (\text{BK-to-REAL2}(P))_2 + n_{23}}{n_{31} \cdot (\text{BK-to-REAL2}(P))_1 + n_{32} \cdot (\text{BK-to-REAL2}(P))_2 + n_{33}}, 1]$. The theorem is a consequence of (33).

- (42) Let us consider an element h of the subgroup of K -isometries, an invertible square matrix N over \mathbb{R}_F of dimension 3, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , and an element u_2 of \mathcal{E}_T^2 . Suppose $h =$ the homography of N and $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $u_2 \in$ the inside of circle(0,0,1). Then $n_{31} \cdot u_2(1) + n_{32} \cdot u_2(2) + n_{33} \neq 0$. The theorem is a consequence of (30).

- (43) Let us consider a positive real number r , and an element u of \mathcal{E}_T^2 . If $u \in$ circle(0,0, r), then u is not zero.

- (44) Let us consider an element h of the subgroup of K -isometries, an invertible square matrix N over \mathbb{R}_F of dimension 3, elements $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ of \mathbb{R}_F , and an element u_2 of \mathcal{E}_T^2 . Suppose $h =$ the homography of N and $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$ and $u_2 \in$ the closed inside of circle(0,0,1). Then $n_{31} \cdot u_2(1) + n_{32} \cdot u_2(2) + n_{33} \neq 0$. The theorem is a consequence of (30), (43), and (32).

- (45) Let us consider real numbers a, b, c, d, e, f, r . Suppose $(1-r) \cdot [a, b, 1] + r \cdot [c, d, 1] = [e, f, 1]$. Then $(1-r) \cdot [a, b] + r \cdot [c, d] = [e, f]$.

- (46) Let us consider points P, Q, R, P', Q', R' of BK-model-Plane, elements p, q, r, p', q', r' of the BK-model, an element h of the subgroup of K -isometries, and an invertible square matrix N over \mathbb{R}_F of dimension 3. Suppose $h =$ the homography of N and Q lies between P and R and $P = p$ and $Q = q$ and $R = r$ and $p' =$ (the homography of N)(p) and $q' =$ (the homography of N)(q) and $r' =$ (the homography of N)(r) and

$P' = p'$ and $Q' = q'$ and $R' = r'$. Then Q' lies between P' and R' .

PROOF: Consider $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ being elements of \mathbb{R}_F such that $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$. Consider u being a non zero element of \mathcal{E}_T^3 such that the direction of $u = p$ and $u(3) = 1$ and $\text{BK-to-REAL2}(p) = [u(1), u(2)]$. Consider v being a non zero element of \mathcal{E}_T^3 such that the direction of $v = r$ and $v(3) = 1$ and $\text{BK-to-REAL2}(r) = [v(1), v(2)]$. Consider w being a non zero element of \mathcal{E}_T^3 such that the direction of $w = q$ and $w(3) = 1$ and $\text{BK-to-REAL2}(q) = [w(1), w(2)]$.

Reconsider $m_1 = n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13}$, $m_2 = n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23}$, $m_3 = n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}$, $m_4 = n_{11} \cdot v(1) + n_{12} \cdot v(2) + n_{13}$, $m_5 = n_{21} \cdot v(1) + n_{22} \cdot v(2) + n_{23}$, $m_6 = n_{31} \cdot v(1) + n_{32} \cdot v(2) + n_{33}$, $m_7 = n_{11} \cdot w(1) + n_{12} \cdot w(2) + n_{13}$, $m_8 = n_{21} \cdot w(1) + n_{22} \cdot w(2) + n_{23}$, $m_9 = n_{31} \cdot w(1) + n_{32} \cdot w(2) + n_{33}$ as a real number. $\text{BKtoT2}(P) = \text{BK-to-REAL2}(p)$ and $\text{BKto}\hat{\text{T}}2(P) = \text{BK-to-REAL2}(p)$ and $\text{BKtoT2}(Q) = \text{BK-to-REAL2}(q)$ and $\text{BKto}\hat{\text{T}}2(Q) = \text{BK-to-REAL2}(q)$ and $\text{BKtoT2}(R) = \text{BK-to-REAL2}(r)$ and $\text{BKto}\hat{\text{T}}2(R) = \text{BK-to-REAL2}(r)$. Consider l being a real number such that $0 \leq l \leq 1$ and $\text{BKto}\hat{\text{T}}2(Q) = (1-l) \cdot (\text{BKto}\hat{\text{T}}2(P)) + l \cdot (\text{BKto}\hat{\text{T}}2(R))$.

Set $r = \frac{l \cdot m_6}{(1-l) \cdot m_3 + l \cdot m_6} \cdot (1-r) \cdot [\frac{m_1}{m_3}, \frac{m_2}{m_3}, 1] + r \cdot [\frac{m_4}{m_6}, \frac{m_5}{m_6}, 1] = [\frac{m_7}{m_9}, \frac{m_8}{m_9}, 1]$. $0 \leq r \leq 1$. $\text{BKtoT2}(P') = \text{BK-to-REAL2}(p')$ and $\text{BKto}\hat{\text{T}}2(P') = \text{BK-to-REAL2}(p')$ and $\text{BKtoT2}(Q') = \text{BK-to-REAL2}(q')$ and $\text{BKto}\hat{\text{T}}2(Q') = \text{BK-to-REAL2}(q')$ and $\text{BKtoT2}(R') = \text{BK-to-REAL2}(r')$ and $\text{BKto}\hat{\text{T}}2(R') = \text{BK-to-REAL2}(r')$. \square

Let P be a point of the projective space over \mathcal{E}_T^3 . We say that P is point at ∞ if and only if

(Def. 4) there exists a non zero element u of \mathcal{E}_T^3 such that $P =$ the direction of u and $(u)_3 = 0$.

Now we state the proposition:

(47) Let us consider a point P of the projective space over \mathcal{E}_T^3 . Suppose there exists a non zero element u of \mathcal{E}_T^3 such that $P =$ the direction of u and $(u)_3 \neq 0$. Then P is not point at ∞ .

Note that there exists a point of the projective space over \mathcal{E}_T^3 which is point at ∞ and there exists a point of the projective space over \mathcal{E}_T^3 which is non point at ∞ .

Let P be a non point at ∞ point of the projective space over \mathcal{E}_T^3 . The functor $\text{RP3toREAL2}(P)$ yielding an element of \mathcal{R}^2 is defined by

(Def. 5) there exists a non zero element u of \mathcal{E}_T^3 such that $P =$ the direction of u and $(u)_3 = 1$ and $it = [(u)_1, (u)_2]$.

The functor $\text{RP3toT2}(P)$ yielding a point of $\text{TarskiEuclid2Space}$ is defined by the term

(Def. 6) $\text{RP3toREAL2}(P)$.

Now we state the propositions:

- (48) Let us consider non point at ∞ elements P, Q, R, P', Q', R' of the projective space over \mathcal{E}_T^3 , an element h of the subgroup of K -isometries, and an invertible square matrix N over \mathbb{R}_F of dimension 3.

Suppose $h =$ the homography of N and $P, Q \in$ the BK-model and $R \in$ the absolute and $P' =$ (the homography of N)(P) and $Q' =$ (the homography of N)(Q) and $R' =$ (the homography of N)(R) and $\text{RP3toT2}(Q)$ lies between $\text{RP3toT2}(P)$ and $\text{RP3toT2}(R)$.

Then $\text{RP3toT2}(Q')$ lies between $\text{RP3toT2}(P')$ and $\text{RP3toT2}(R')$.

PROOF: Consider $n_{11}, n_{12}, n_{13}, n_{21}, n_{22}, n_{23}, n_{31}, n_{32}, n_{33}$ being elements of \mathbb{R}_F such that $N = \langle \langle n_{11}, n_{12}, n_{13} \rangle, \langle n_{21}, n_{22}, n_{23} \rangle, \langle n_{31}, n_{32}, n_{33} \rangle \rangle$. Consider u being a non zero element of \mathcal{E}_T^3 such that $P =$ the direction of u and $(u)_3 = 1$ and $\text{RP3toREAL2}(P) = [(u)_1, (u)_2]$. Consider v being a non zero element of \mathcal{E}_T^3 such that $R =$ the direction of v and $(v)_3 = 1$ and $\text{RP3toREAL2}(R) = [(v)_1, (v)_2]$. Consider w being a non zero element of \mathcal{E}_T^3 such that $Q =$ the direction of w and $(w)_3 = 1$ and $\text{RP3toREAL2}(Q) = [(w)_1, (w)_2]$.

Reconsider $m_1 = n_{11} \cdot u(1) + n_{12} \cdot u(2) + n_{13}, m_2 = n_{21} \cdot u(1) + n_{22} \cdot u(2) + n_{23}, m_3 = n_{31} \cdot u(1) + n_{32} \cdot u(2) + n_{33}, m_4 = n_{11} \cdot v(1) + n_{12} \cdot v(2) + n_{13}, m_5 = n_{21} \cdot v(1) + n_{22} \cdot v(2) + n_{23}, m_6 = n_{31} \cdot v(1) + n_{32} \cdot v(2) + n_{33}, m_7 = n_{11} \cdot w(1) + n_{12} \cdot w(2) + n_{13}, m_8 = n_{21} \cdot w(1) + n_{22} \cdot w(2) + n_{23}, m_9 = n_{31} \cdot w(1) + n_{32} \cdot w(2) + n_{33}$ as a real number.

Consider l being a real number such that $0 \leq l \leq 1$ and $\text{RP3to}\hat{\text{T}}2(Q) = (1-l) \cdot (\text{RP3to}\hat{\text{T}}2(P)) + l \cdot (\text{RP3to}\hat{\text{T}}2(R))$. Set $r = \frac{l \cdot m_6}{(1-l) \cdot m_3 + l \cdot m_6} \cdot (1-r) \cdot [\frac{m_1}{m_3}, \frac{m_2}{m_3}, 1] + r \cdot [\frac{m_4}{m_6}, \frac{m_5}{m_6}, 1] = [\frac{m_7}{m_9}, \frac{m_8}{m_9}, 1]$. $0 \leq r \leq 1$. \square

- (49) Let us consider real numbers a, b, c , and elements u, v, w of \mathcal{E}_T^3 . Suppose $a \neq 0$ and $a + b + c = 0$ and $a \cdot u + b \cdot v + c \cdot w = 0_{\mathcal{E}_T^3}$. Then $u \in \text{Line}(v, w)$.
- (50) Let us consider non point at ∞ points P, Q, R of the projective space over \mathcal{E}_T^3 , and non zero elements u, v, w of \mathcal{E}_T^3 . Suppose $P =$ the direction of u and $Q =$ the direction of v and $R =$ the direction of w and $(u)_3 = 1$ and $(v)_3 = 1$ and $(w)_3 = 1$. Then P, Q and R are collinear if and only if u, v and w are collinear. The theorem is a consequence of (49).
- (51) Let us consider elements u, v, w of \mathcal{E}_T^3 . Suppose $u \in \mathcal{L}(v, w)$. Then $[(u)_1, (u)_2] \in \mathcal{L}([(v)_1, (v)_2], [(w)_1, (w)_2])$.
- (52) Let us consider elements u, v, w of \mathcal{E}_T^2 . Suppose $u \in \mathcal{L}(v, w)$. Then $[(u)_1, (u)_2, 1] \in \mathcal{L}([(v)_1, (v)_2, 1], [(w)_1, (w)_2, 1])$.

PROOF: Consider r being a real number such that $0 \leq r$ and $r \leq 1$ and $u = (1 - r) \cdot v + r \cdot w$. Reconsider $u' = [(u)_1, (u)_2, 1]$, $v' = [(v)_1, (v)_2, 1]$, $w' = [(w)_1, (w)_2, 1]$ as an element of \mathcal{E}_T^3 . $u' = (1 - r) \cdot v' + r \cdot w'$. \square

- (53) Let us consider non point at ∞ points P, Q, R of the projective space over \mathcal{E}_T^3 . Then P, Q and R are collinear if and only if $\text{RP3toT2}(P)$, $\text{RP3toT2}(Q)$ and $\text{RP3toT2}(R)$ are collinear. The theorem is a consequence of (50), (51), and (52).
- (54) Let us consider elements u, v, w of \mathcal{E}_T^2 . Suppose u, v and w are collinear. Then $[(u)_1, (u)_2, 1]$, $[(v)_1, (v)_2, 1]$ and $[(w)_1, (w)_2, 1]$ are collinear. The theorem is a consequence of (52).
- (55) Let us consider non point at ∞ elements P, Q, P_1 of the projective space over \mathcal{E}_T^3 . Suppose $P, Q \in$ the BK-model and $P_1 \in$ the absolute. Then $\text{RP3toT2}(P_1)$ does not lie between $\text{RP3toT2}(Q)$ and $\text{RP3toT2}(P)$. The theorem is a consequence of (52) and (27).

The functor Dir001 yielding a non point at ∞ element of the projective space over \mathcal{E}_T^3 is defined by the term

(Def. 7) the direction of $[0, 0, 1]$.

The functor Dir101 yielding a non point at ∞ element of the projective space over \mathcal{E}_T^3 is defined by the term

(Def. 8) the direction of $[1, 0, 1]$.

Now we state the propositions:

- (56) Let us consider non point at ∞ elements P, Q of the projective space over \mathcal{E}_T^3 . Suppose $P, Q \in$ the absolute. Then $\overline{\text{RP3toT2}(\text{Dir001}) \text{RP3toT2}(P)} \cong \overline{\text{RP3toT2}(\text{Dir101}) \text{RP3toT2}(Q)}$.
- (57) Let us consider non point at ∞ elements P, Q, R of the projective space over \mathcal{E}_T^3 , and non zero elements u, v, w of \mathcal{E}_T^3 . Suppose $P, Q \in$ the absolute and $P \neq Q$ and $P =$ the direction of u and $Q =$ the direction of v and $R =$ the direction of w and $(u)_3 = 1$ and $(v)_3 = 1$ and $w = [\frac{(u)_1 + (v)_1}{2}, \frac{(u)_2 + (v)_2}{2}, 1]$. Then $R \in$ the BK-model.

PROOF: Reconsider $u' = [u(1), u(2)]$, $v' = [v(1), v(2)]$ as an element of \mathcal{E}_T^2 . $u' \neq v'$. Reconsider $r_8 = [(w)_1, (w)_2]$ as an element of the inside of circle(0,0,1). Consider R_3 being an element of \mathcal{E}_T^2 such that $R_3 = r_8$ and $\text{REAL2-to-BK}(r_8) =$ the direction of $[(R_3)_1, (R_3)_2, 1]$. \square

- (58) Let us consider points R_1, R_2 of TarskiEuclid2Space. Suppose $\hat{R}_1, \hat{R}_2 \in$ circle(0,0,1) and $R_1 \neq R_2$. Then there exists an element P of BK-model-Plane such that $\text{BKtoT2}(P)$ lies between R_1 and R_2 . The theorem is a consequence of (47), (57), and (26).
- (59) Let us consider non point at ∞ elements P, Q of the projective space

over \mathcal{E}_T^3 . If $\text{RP3toT2}(P) = \text{RP3toT2}(Q)$, then $P = Q$.

- (60) Let us consider non point at ∞ elements R_1, R_2 of the projective space over \mathcal{E}_T^3 . Suppose $R_1, R_2 \in$ the absolute and $R_1 \neq R_2$. Then there exists an element P of BK-model-Plane such that $\text{BKtoT2}(P)$ lies between $\text{RP3toT2}(R_1)$ and $\text{RP3toT2}(R_2)$. The theorem is a consequence of (59) and (58).
- (61) Let us consider points P, Q, R of TarskiEuclid2Space. Suppose Q lies between P and R and $\hat{P}, \hat{R} \in$ the inside of circle(0,0,1). Then $\hat{Q} \in$ the inside of circle(0,0,1).

Let us consider a non point at ∞ element P of the projective space over \mathcal{E}_T^3 .

- (62) If $P \in$ the absolute, then $\text{RP3toREAL2}(P) \in$ circle(0,0,1).
- (63) If $P \in$ the BK-model, then $\text{RP3toREAL2}(P) \in$ the inside of circle(0,0,1). The theorem is a consequence of (26).
- (64) Let us consider a non point at ∞ point P of the projective space over \mathcal{E}_T^3 , and an element Q of the BK-model. If $P = Q$, then $\text{RP3toREAL2}(P) = \text{BK-to-REAL2}(Q)$. The theorem is a consequence of (26).
- (65) Let us consider non point at ∞ elements P, Q, R_1, R_2 of the projective space over \mathcal{E}_T^3 . Suppose $P \neq Q$ and $P \in$ the BK-model and $R_1, R_2 \in$ the absolute and $\text{RP3toT2}(Q)$ lies between $\text{RP3toT2}(P)$ and $\text{RP3toT2}(R_1)$ and $\text{RP3toT2}(Q)$ lies between $\text{RP3toT2}(P)$ and $\text{RP3toT2}(R_2)$. Then $R_1 = R_2$. The theorem is a consequence of (60), (59), (62), (64), (8), and (61).
- (66) Let us consider non point at ∞ elements P, Q, P_1, P_2 of the projective space over \mathcal{E}_T^3 . Suppose $P \neq Q$ and $P, Q \in$ the BK-model and $P_1, P_2 \in$ the absolute and $P_1 \neq P_2$ and P, Q and P_1 are collinear and P, Q and P_2 are collinear. Then
- (i) $\text{RP3toT2}(P)$ lies between $\text{RP3toT2}(Q)$ and $\text{RP3toT2}(P_1)$, or
 - (ii) $\text{RP3toT2}(P)$ lies between $\text{RP3toT2}(Q)$ and $\text{RP3toT2}(P_2)$.

The theorem is a consequence of (55), (53), and (65).

Let us consider elements P, Q of the BK-model. Now we state the propositions:

- (67) Suppose $P \neq Q$. Then there exists an element R of the absolute such that for every non point at ∞ elements p, q, r of the projective space over \mathcal{E}_T^3 such that $p = P$ and $q = Q$ and $r = R$ holds $\text{RP3toT2}(p)$ lies between $\text{RP3toT2}(q)$ and $\text{RP3toT2}(r)$. The theorem is a consequence of (47) and (66).
- (68) Suppose $P \neq Q$. Then there exists an element R of the absolute such that for every non point at ∞ elements p, q, r of the projective space over

\mathcal{E}_T^3 such that $p = P$ and $q = Q$ and $r = R$ holds $\text{RP3toT2}(q)$ lies between $\text{RP3toT2}(p)$ and $\text{RP3toT2}(r)$. The theorem is a consequence of (67).

- (69) The direction of $[1, 0, 1]$ is an element of the absolute.
- (70) Let us consider points a, b of BK-model-Plane. Then $\overline{aa} \cong \overline{bb}$. The theorem is a consequence of (69).
- (71) Every element of the BK-model is a non point at ∞ element of the projective space over \mathcal{E}_T^3 . The theorem is a consequence of (47).
- (72) Every element of the absolute is a non point at ∞ element of the projective space over \mathcal{E}_T^3 . The theorem is a consequence of (47).
- (73) Let us consider an element P of the BK-model, and a non point at ∞ element P' of the projective space over \mathcal{E}_T^3 . If $P = P'$, then $\text{RP3toREAL2}(P') = \text{BK-to-REAL2}(P)$. The theorem is a consequence of (26).
- (74) Let us consider points a, q, b, c of BK-model-Plane. Then there exists a point x of BK-model-Plane such that
- (i) a lies between q and x , and
 - (ii) $\overline{ax} \cong \overline{bc}$.

The theorem is a consequence of (71), (68), (72), (12), (70), (48), and (73).

- (75) Let us consider points P, Q of BK-model-Plane.
If $\text{BKtoT2}(P) = \text{BKtoT2}(Q)$, then $P = Q$.
- (76) Let us consider real numbers a, b, r , and elements P, Q, R of \mathcal{E}_T^2 . Suppose $P, R \in$ the inside of circle(a, b, r). Then $\mathcal{L}(P, R) \subseteq$ the inside of circle(a, b, r).

2. THE AXIOM OF SEGMENT CONSTRUCTION

Now we state the proposition:

- (77) BK-model-Plane satisfies the axiom of segment construction.

3. THE AXIOM OF BETWEENNESS IDENTITY

Now we state the proposition:

- (78) BK-model-Plane satisfies the axiom of betweenness identity. The theorem is a consequence of (12) and (75).

4. THE AXIOM OF PASCH

Now we state the proposition:

- (79) BK-model-Plane satisfies the axiom of Pasch. The theorem is a consequence of (12), (8), (25), and (10).

REFERENCES

- [1] Grzegorz Bancerek, Czesław Byliński, Adam Grabowski, Artur Korniłowicz, Roman Matuszewski, Adam Naumowicz, Karol Pąk, and Josef Urban. Mizar: State-of-the-art and beyond. In Manfred Kerber, Jacques Carette, Cezary Kaliszyk, Florian Rabe, and Volker Sorge, editors, *Intelligent Computer Mathematics*, volume 9150 of *Lecture Notes in Computer Science*, pages 261–279. Springer International Publishing, 2015. ISBN 978-3-319-20614-1. doi:10.1007/978-3-319-20615-8_17.
- [2] Eugenio Beltrami. Saggio di interpretazione della geometria non-euclidea. *Giornale di Matematiche*, 6:284–322, 1868.
- [3] Eugenio Beltrami. Essai d’interprétation de la géométrie non-euclidéenne. In *Annales scientifiques de l’École Normale Supérieure. Trad. par J. Hoüel*, volume 6, pages 251–288. Elsevier, 1869.
- [4] Karol Borsuk and Wanda Szmielew. *Foundations of Geometry*. North Holland, 1960.
- [5] Karol Borsuk and Wanda Szmielew. *Podstawy geometrii*. Państwowe Wydawnictwo Naukowe, Warszawa, 1955 (in Polish).
- [6] Roland Coghetto. Homography in \mathbb{RP}^2 . *Formalized Mathematics*, 24(4):239–251, 2016. doi:10.1515/forma-2016-0020.
- [7] Roland Coghetto. Klein-Beltrami model. Part I. *Formalized Mathematics*, 26(1):21–32, 2018. doi:10.2478/forma-2018-0003.
- [8] Roland Coghetto. Klein-Beltrami model. Part III. *Formalized Mathematics*, 28(1):1–7, 2020. doi:10.2478/forma-2020-0001.
- [9] Kanchun, Hiroshi Yamazaki, and Yatsuka Nakamura. Cross products and tripple vector products in 3-dimensional Euclidean space. *Formalized Mathematics*, 11(4):381–383, 2003.
- [10] Timothy James McKenzie Makarios. A mechanical verification of the independence of Tarski’s Euclidean Axiom. Victoria University of Wellington, New Zealand, 2012. Master’s thesis.
- [11] William Richter, Adam Grabowski, and Jesse Alama. Tarski geometry axioms. *Formalized Mathematics*, 22(2):167–176, 2014. doi:10.2478/forma-2014-0017.

Accepted December 30, 2019
